

PAPER – II
MATHEMATICAL SCIENCES

Note : Attempt all the questions. Each question carries *two* (2) marks.

1. Composite number n is _____.
- 1) a prime number and $n > 1$
 - 2) non-prime number and $n < 1$
 - 3) non-prime number and $n > 1$
 - 4) a prime number and $n < 1$
2. A function $f(x)$ has no jump discontinuity at $x = a$ if _____.
- 1) $f(a+) = f(a-) = f(a)$
 - 2) $f(a+) \neq f(a-)$
 - 3) $f(a+) \neq f(a)$
 - 4) $f(a+) = f(a-) \neq f(a)$
3. A subset S of a vector space V satisfying $V = L(S)$ is a basis if _____.
- 1) S is linearly dependent
 - 2) S is linearly independent
 - 3) V is a field
 - 4) S is a field
4. The value of the determinant $\begin{vmatrix} b^2c^2 & bc & b+c \\ c^2a^2 & ca & c+a \\ a^2b^2 & ab & a+b \end{vmatrix}$ is
- 1) abc
 - 2) $a^2b^2c^2$
 - 3) $bc + ca + ab$
 - 4) zero
5. If $\dim W = m$, $\dim V = n$ and $W \subset V$ then $\dim(V/W)$ is _____.
- 1) $m + n$
 - 2) $n - m$
 - 3) $m - n$
 - 4) mn

6. The value of $\lim_{z \rightarrow 0} \frac{\bar{z}}{z}$ is
- 1) 0
 - 2) 1
 - 3) $1/2$
 - 4) Limit does not exist
7. The fixed points of the bilinear transformation $w = \frac{z}{2-z}$ are
- 1) 0, 0
 - 2) 0, 1
 - 3) 0, $1/2$
 - 4) 1, $1/2$
8. The primitive roots *modulo* 19 is _____.
- 1) 18
 - 2) 6
 - 3) 5
 - 4) 12
9. In the ring of even integers $2Z$, the ideal $I = \langle 4 \rangle$ is
- 1) Integral domain
 - 2) Principal ideal
 - 3) Maximal but not prime
 - 4) Maximal and prime
10. If D is an integral domain and $D[x]$ is a principal ideal domain, then D becomes a
- 1) Ring
 - 2) Field
 - 3) Integral domain
 - 4) Ideal

11. Any infinite subset A of a discrete topological space X is

- 1) compact
- 2) locally compact
- 3) not compact
- 4) sequentially compact

12. The general solution of $\frac{dy}{dx} = \frac{y}{x} + \frac{1}{\sin\left(\frac{y}{x}\right)}$ is

- 1) $y = x \operatorname{arc} \cos(x - c)$
- 2) $y = x \operatorname{arc} \sin(x - c)$
- 3) $y = x \operatorname{arc} \tan(x - c)$
- 4) $y = x \sin(x - c)$

13. Which of the following is elliptic?

- 1) Laplace equation
- 2) Wave equation
- 3) Heat equation
- 4) $u_{xx} + 2u_{xy} - 4u_{yy} = 0$

14. The complete integral of the PDE $pq = 1$ is

- 1) $z = ax + 1/a y + c$
- 2) $z = ax + 1/a y$
- 3) $z = ax + y + c$
- 4) $z = x + 1/a y + c$

15. For the fastest rate of convergence of the method $x_{n-1} = \frac{\alpha x_n - x_n^2 + 1}{\alpha + x_n}$. The value of α is

(Given α is exact root)

1) $\alpha = \frac{1}{\alpha^2}$

2) $\alpha = \frac{2}{\alpha^3}$

3) $\alpha = \frac{1}{\alpha^3}$

4) $\alpha = \frac{3}{\alpha^2}$

16. If Euler's characteristic equation $\frac{\partial F}{\partial y} - \frac{d}{dx} \left(\frac{\partial F}{\partial y'} \right) = 0$ vanishes identically, then the indefinite integral $\int F(x, y, y') dx$ can be evaluated as a function of

1) x only

2) x and y

3) y only

4) F, x and y

17. The integral I has strong minimum if

1) The arc AB of the arc of the integration Γ_e , contains no point conjugate to either A or B

2) The arc AB of the arc of integration Γ_e , contains point conjugate to either A or B

3) The arc AB of the arc of integration Γ_e , contains point conjugate to both A or B

4) The arc AB of the arc of the integration Γ_e , contains no point conjugate to neither A nor B

18. Degree of freedom is defined as

- 1) The minimum number of independent coordinates required to specify the system
- 2) The maximum number of independent coordinates required to specify the system
- 3) The minimum number of dependent coordinates required to specify the system
- 4) The maximum number of dependent coordinates required to specify the system

19. Non-holonomic constraints are

- 1) The constraints that can be expressed as equation form
- 2) The constraints that cannot be expressed as equation form
- 3) Equation of constraints that contain time as explicit variable
- 4) Equation of constraints that does not contain time as explicit variable

20. Lagrange's bracket is

- 1) Canonical invariant
- 2) Canonical variant
- 3) Non-invariant
- 4) Euler's invariant

21. Two dice are thrown. Find the probability that the total of the numbers on the top face is 9.

- 1) $\frac{3}{9}$
- 2) $\frac{4}{9}$
- 3) $\frac{5}{36}$
- 4) $\frac{4}{36}$

22. A non-null persistent and a periodic state is called

- 1) Regular
- 2) Irregular
- 3) Ergodic
- 4) Non-Ergodic

23. The variance of maximum likelihood estimate for the parameter λ of a Poisson distribution on the basis of a Poisson distribution on the basis of a sample of size n is

- 1) λ
- 2) $\frac{n}{\lambda}$
- 3) $\frac{\lambda}{n}$
- 4) $\frac{\lambda}{n^2}$

24. Buses arrive for cleaning at a central depot in groups of five every hour on the hour. The buses are serviced in random order, one at a time. Each bus requires 11 min to service completely and it leaves the depot as soon as it is clean. Then the average number of buses in the depot is

- 1) 2
- 2) 2.5
- 3) 2.75
- 4) 3

25. Subset of R which is a neighborhood of 3 is

- | | |
|-----------|-----------|
| 1) [3, 6] | 2) [3, 6) |
| 3) (2, 4) | 4) (3, 6) |

26. The series $\sum_{r=1}^{\alpha} (-1)^{r-1}$

- 1) Oscillates finitely
- 2) Divergent
- 3) Convergent
- 4) Oscillates infinitely

27. The sequence $\left[\frac{\cos \frac{n\pi}{2}}{n} \right]_{n=1}^{\infty}$ is _____.

- 1) convergent to 0
- 2) divergent
- 3) convergent to 1
- 4) convergent to $\frac{1}{2}$

28. The radius of convergence of the series $\frac{x + .2}{1} + \frac{(x + .2)^2}{2} + \dots + \frac{(x + .2)^n}{n} + \dots$

- | | |
|------|------------------|
| 1) 1 | 2) ∞ |
| 3) 0 | 4) $\frac{1}{2}$ |

29. The product of two orthogonal matrices is orthogonal and that the inverse of an orthogonal matrix is
- 1) Symmetric
 - 2) Orthogonal
 - 3) Skew-symmetric
 - 4) Hermitian
30. Let V be an n -dimensional vector space and let $T : V \rightarrow V$ be a linear map such that $\text{null}(T) = \text{range}(T)$. Then
- 1) n is odd
 - 2) n is even
 - 3) n is neither odd or even
 - 4) n is not defined
31. Every square matrix satisfies its own characteristic equation. This is
- 1) Cauchy's theorem
 - 2) Cayley-Hamilton theorem
 - 3) Eigen value theorem
 - 4) Sylow's theorem
32. A function $f(z) = \text{Re}(z)$ is
- 1) analytic
 - 2) nowhere differentiable
 - 3) continuous
 - 4) discontinuous
33. The real part of $\exp(\exp i\theta)$ is
- 1) $e^{\cos \theta}$
 - 2) $e^{\cos \theta} \sin(\sin \theta)$
 - 3) $e^{\cos \theta} \cos(\cos \theta)$
 - 4) $e^{\cos \theta} \cos(\sin \theta)$

34. If $\pi(N)$ denotes the number of prime numbers less than or equal to N then $\pi(6) =$
- 1) 2
 - 2) 5
 - 3) 1
 - 4) 4
35. Which of the following statement is wrong?
- 1) Every subspace of discrete space is also discrete
 - 2) Every subspace of an indiscrete space is indiscrete
 - 3) Every non-empty open subset of an indiscrete space X is dense in X
 - 4) Every non-empty open subset of an indiscrete space X is not dense in X
36. Let $X = N$ be equipped with the topology generated by the basis consisting of sets $A_n = \{n, n + 1, n + 2 \dots\}$, $n \in N$ then X is
- 1) Compact and connected
 - 2) Hausdorff and compact
 - 3) Hausdorff and connected
 - 4) Neither compact nor connected
37. Every convergent sequence in a Hausdorff space has
- 1) exactly two different limit points
 - 2) no limit point
 - 3) a unique limit point
 - 4) more than one limit point
38. Let X be a topological space with finitely many connected components: Then each connected components is
- 1) closed in X
 - 2) open in X
 - 3) neither open nor closed in X
 - 4) both open and closed in X

39. The PDE $(1 + x^2)u_{xx} + (1 + y^2)u_{yy} + xu_x + yu_y = 0$ is of type
- 1) Parabolic
 - 2) Elliptic
 - 3) Hyperbolic
 - 4) Laplace
40. The PDE $x^2(y - 1)z_{xx} - x(y^2 - 1)z_{xy} + y(y^2 - 1)z_{yy} + z_x = 0$ is hyperbolic in the entire xy -plane except along
- 1) x -axis
 - 2) y -axis
 - 3) a line parallel to y axis
 - 4) a line parallel to x axis
41. In Newton-Cotes formula, if $f(x)$ is interpolated at equally spaced nodes by a polynomial of degree one, then it represents _____.
- 1) Trapezoidal rule
 - 2) Simpson rule
 - 3) Three-eight rule
 - 4) Booles rule
42. By Newton's method $f(x) = x^5 - x^3 + 3$ and if $x_n = 1$ then x_{n+1} is
- 1) $-\frac{1}{2}$
 - 2) $\frac{1}{2}$
 - 3) $\frac{3}{2}$
 - 4) $-\frac{3}{2}$

43. The solution of Fredholm integral equation $y(x) = x + e^x - \int_0^1 xt y(t) dt$

- 1) $y(x) = e^x$
- 2) $y(x) = e^{-x}$
- 3) $y(x) = x$
- 4) $y(x) = x + 1$

44. Consider the Fredholm integral equation of the first kind $\frac{1}{4}e^x = \int_0^{\frac{1}{4}} e^{x-t} y(t) dt$. The solution is

- 1) $y(x) = e^x$
- 2) $y(x) = \cos x$
- 3) $y(x) = \sin x$
- 4) $y(x) = e^{-x}$

45. Any solution of homogeneous Volterra integral equation of the second kind $\phi(x) - \lambda \int_0^x K(x \cdot y) \phi(x) dy = 0$ in L_2 -space is

- 1) Necessarily a zero function
- 2) Necessarily a non-zero function
- 3) Absolute function
- 4) Constant function

46. The eigen value I of the following Fredholm integral equation $y(x) = I \int_0^1 x^2 t y(t) dt$ is

- 1) -2
- 2) 2
- 3) 4
- 4) -4

47. From a pack of 52 cards, one card is drawn at random. Find the probability of getting a queen.
- 1) $\frac{1}{13}$
 - 2) $\frac{2}{13}$
 - 3) $\frac{10}{21}$
 - 4) $\frac{12}{21}$
48. Poisson distribution is a limiting case of
- 1) Uniform distribution
 - 2) Exponential distribution
 - 3) Geometric distribution
 - 4) Binomial distribution
49. If X is uniformly distributed over $(0, 10)$ find $P(X < 2)$
- 1) $\frac{3}{5}$
 - 2) $\frac{2}{5}$
 - 3) $\frac{1}{5}$
 - 4) $\frac{4}{5}$
50. If the one-step transition probability does not depend on the step (ie., $p_{ij}(n-1, n) = p_{ij}(m-1, m)$) the Markov chain is called a
- 1) Non-homogeneous
 - 2) Homogeneous
 - 3) Irreducible
 - 4) Reducible

ROUGH WORK

ROUGH WORK