


## Problems and Solutions: INMO-2015

1. Let  $ABC$  be a right-angled triangle with  $\angle B = 90^\circ$ . Let  $BD$  be the altitude from  $B$  on to  $AC$ . Let  $P, Q$  and  $I$  be the incentres of triangles  $ABD, CBD$  and  $ABC$  respectively. Show that the circumcentre of of the triangle  $PIQ$  lies on the hypotenuse  $AC$ .

**Solution:** We begin with the following lemma:

**Lemma:** Let  $XYZ$  be a triangle with  $\angle XYZ = 90 + \alpha$ . Construct an isosceles triangle  $XEZ$ , externally on the side  $XZ$ , with base angle  $\alpha$ . Then  $E$  is the circumcentre of  $\triangle XYZ$ .

**Proof of the Lemma:** Draw  $ED \perp XZ$ . Then  $DE$  is the perpendicular bisector of  $XZ$ . We also observe that  $\angle XED = \angle ZED = 90 - \alpha$ . Observe that  $E$  is on the perpendicular bisector of  $XZ$ . Construct the circumcircle of  $XYZ$ . Draw perpendicular bisector of  $XY$  and let it meet  $DE$  in  $F$ . Then  $F$  is the circumcentre of  $\triangle XYZ$ . Join  $XF$ . Then  $\angle XFD = 90 - \alpha$ . But we know that  $\angle XED = 90 - \alpha$ . Hence  $E = F$ .




Let  $r_1, r_2$  and  $r$  be the inradii of the triangles  $ABD, CBD$  and  $ABC$  respectively. Join  $PD$  and  $DQ$ . Observe that  $\angle PDQ = 90^\circ$ . Hence

$$PQ^2 = PD^2 + DQ^2 = 2r_1^2 + 2r_2^2.$$

Let  $s_1 = (AB + BD + DA)/2$ . Observe that  $BD = ca/b$  and  $AD = \sqrt{AB^2 - BD^2} = \sqrt{c^2 - (ca/b)^2} = c^2/b$ . This gives  $s_1 = cs/b$ . But  $r_1 = s_1 - c = (c/b)(s - b) = cr/b$ . Similarly,  $r_2 = ar/b$ . Hence

$$PQ^2 = 2r^2 \left( \frac{c^2 + a^2}{b^2} \right) = 2r^2.$$

Consider  $\triangle PIQ$ . Observe that  $\angle PIQ = 90 + (B/2) = 135$ . Hence  $PQ$  subtends  $90^\circ$  on the circumference of the circumcircle of  $\triangle PIQ$ . But we have seen that  $\angle PDQ = 90^\circ$ . Now construct a circle with  $PQ$  as diameter. Let it cut  $AC$  again in  $K$ . It follows that  $\angle PKQ = 90^\circ$  and the points  $P, D, K, Q$  are concyclic. We also notice  $\angle KPQ = \angle KDQ = 45^\circ$  and  $\angle PQK = \angle PDK = 45^\circ$ .





length of the non-periodic part of the infinite decimal expansion of  $1/n$ .

Write

$$\frac{1}{n} = 0.a_1a_2 \cdots a_r \overline{b_1b_2 \cdots b_s}.$$

We show that  $r = \max(\nu_2(n), \nu_5(n))$ .

Let  $a$  and  $b$  be the numbers  $a_1a_2 \cdots a_r$  and  $b = b_1b_2 \cdots b_s$  respectively. (Here  $a_1$  and  $b_1$  can be both 0.) Then

$$\frac{1}{n} = \frac{1}{10^r} \left( a + \sum_{k \geq 1} \frac{b}{(10^s)^k} \right) = \frac{1}{10^r} \left( a + \frac{b}{10^s - 1} \right).$$

Thus we get  $10^r(10^s - 1) = n((10^s - 1)a + b)$ . It shows that  $r \geq \max(\nu_2(n), \nu_5(n))$ . Suppose  $r > \max(\nu_2(n), \nu_5(n))$ . Then 10 divides  $b - a$ . Hence the last digits of  $a$  and  $b$  are equal:  $a_r = b_s$ . This means

$$\frac{1}{n} = 0.a_1a_2 \cdots a_{r-1} \overline{b_sb_1b_2 \cdots b_{s-1}}.$$

This contradicts the definition of  $r$ . Therefore  $r = \max(\nu_2(n), \nu_5(n))$ .

3. Find all real functions  $f$  from  $\mathbb{R} \rightarrow \mathbb{R}$  satisfying the relation

$$f(x^2 + yf(x)) = xf(x + y).$$

**Solution:** Put  $x = 0$  and we get  $f(yf(0)) = 0$ . If  $f(0) \neq 0$ , then  $yf(0)$  takes all real values when  $y$  varies over real line. We get  $f(x) \equiv 0$ . Suppose  $f(0) = 0$ . Taking  $y = -x$ , we get  $f(x^2 - xf(x)) = 0$  for all real  $x$ .

Suppose there exists  $x_0 \neq 0$  in  $\mathbb{R}$  such that  $f(x_0) = 0$ . Putting  $x = x_0$  in the given relation we get

$$f(x_0^2) = x_0f(x_0 + y),$$

for all  $y \in \mathbb{R}$ . Now the left side is a constant and hence it follows that  $f$  is a constant function. But the only constant function which satisfies the equation is identically zero function, which is already obtained. Hence we may consider the case where  $f(x) \neq 0$  for all  $x \neq 0$ .

Since  $f(x^2 - xf(x)) = 0$ , we conclude that  $x^2 - xf(x) = 0$  for all  $x \neq 0$ . This implies that  $f(x) = x$  for all  $x \neq 0$ . Since  $f(0) = 0$ , we conclude that  $f(x) = x$  for all  $x \in \mathbb{R}$ .

Thus we have two functions:  $f(x) \equiv 0$  and  $f(x) = x$  for all  $x \in \mathbb{R}$ .

4. There are four basket-ball players  $A, B, C, D$ . Initially, the ball is with  $A$ . The ball is always passed from one person to a different person. In how many ways can the ball come back to  $A$  after **seven** passes? (For example  $A \rightarrow C \rightarrow B \rightarrow D \rightarrow A \rightarrow B \rightarrow C \rightarrow A$  and

$A \rightarrow D \rightarrow A \rightarrow D \rightarrow C \rightarrow A \rightarrow B \rightarrow A$  are two ways in which the ball can come back to  $A$  after seven passes.)

**Solution:** Let  $x_n$  be the number of ways in which  $A$  can get back the ball after  $n$  passes. Let  $y_n$  be the number of ways in which the ball goes back to a fixed person other than  $A$  after  $n$  passes. Then

$$x_n = 3y_{n-1},$$

and

$$y_n = x_{n-1} + 2y_{n-1}.$$

We also have  $x_1 = 0$ ,  $x_2 = 3$ ,  $y_1 = 1$  and  $y_2 = 2$ .

Eliminating  $y_n$  and  $y_{n-1}$ , we get  $x_{n+1} = 3x_{n-1} + 2x_n$ . Thus

$$\begin{aligned} x_3 &= 3x_1 + 2x_2 = 2 \times 3 = 6; \\ x_4 &= 3x_2 + 2x_3 = (3 \times 3) + (2 \times 6) = 9 + 12 = 21; \\ x_5 &= 3x_3 + 2x_4 = (3 \times 6) + (2 \times 21) = 18 + 42 = 60; \\ x_6 &= 3x_4 + 2x_5 = (3 \times 21) + (2 \times 60) = 63 + 120 = 183; \\ x_7 &= 3x_5 + 2x_6 = (3 \times 60) + (2 \times 183) = 180 + 366 = 546. \end{aligned}$$

**Alternate solution:** Since the ball goes back to one of the other 3 persons, we have

$$x_n + 3y_n = 3^n,$$

since there are  $3^n$  ways of passing the ball in  $n$  passes. Using  $x_n = 3y_{n-1}$ , we obtain

$$x_{n-1} + x_n = 3^{n-1},$$


with  $x_1 = 0$ . Thus

$$\begin{aligned} x_7 &= 3^6 - x_6 = 3^6 - 3^5 + x_5 = 3^6 - 3^5 + 3^4 - x_4 = 3^6 - 3^5 + 3^4 - 3^3 + x_3 \\ &= 3^6 - 3^5 + 3^4 - 3^3 + 3^2 - x_2 = 3^6 - 3^5 + 3^4 - 3^3 + 3^2 - 3 \\ &= (2 \times 3^5) + (2 \times 3^3) + (2 \times 3) = 486 + 54 + 6 = 546. \end{aligned}$$

5. Let  $ABCD$  be a convex quadrilateral. Let the diagonals  $AC$  and  $BD$  intersect in  $P$ . Let  $PE$ ,  $PF$ ,  $PG$  and  $PH$  be the altitudes from  $P$  on to the sides  $AB$ ,  $BC$ ,  $CD$  and  $DA$  respectively. Show that  $ABCD$  has an incircle if and only if

$$\frac{1}{PE} + \frac{1}{PG} = \frac{1}{PF} + \frac{1}{PH}.$$

**Solution:** Let  $AP = p$ ,  $BP = q$ ,  $CP = r$ ,  $DP = s$ ;  $AB = a$ ,  $BC = b$ ,  $CD = c$  and  $DA = d$ . Let  $\angle APB = \angle CPD = \theta$ . Then  $\angle BPC = \angle DPA = \pi - \theta$ . Let us also write  $PE = h_1$ ,  $PF = h_2$ ,  $PG = h_3$  and  $PH = h_4$ .



Observe that

$$h_1 a = pq \sin \theta, \quad h_2 b = qr \sin \theta, \quad h_3 c = rs \sin \theta, \quad h_4 d = sp \sin \theta.$$

Hence

$$\frac{1}{h_1} + \frac{1}{h_3} = \frac{1}{h_2} + \frac{1}{h_4}.$$

is equivalent to

$$\frac{a}{pq} + \frac{c}{rs} = \frac{b}{qr} + \frac{d}{sp}.$$

This is the same as

$$ars + cpq = bsp + dqr.$$

Thus we have to prove that  $a+c = b+d$  if and only if  $ars+cpq = bsp+dqr$ .

Now we can write  $a + c = b + d$  as

$$a^2 + c^2 + 2ac = b^2 + d^2 + 2bd.$$

But we know that

$$\begin{aligned} a^2 &= p^2 + q^2 - 2pq \cos \theta, & c^2 &= r^2 + s^2 - 2rs \cos \theta \\ b^2 &= q^2 + r^2 + 2qr \cos \theta, & d^2 &= p^2 + s^2 + 2ps \cos \theta, \end{aligned}$$

Hence  $a + c = b + d$  is equivalent to

$$-pq \cos \theta + -rs \cos \theta + ac = ps \cos \theta + qr \cos \theta + bd.$$

Similarly, by squaring  $ars + cpq = bsp + dqr$  we can show that it is equivalent to

$$-pq \cos \theta + -rs \cos \theta + ac = ps \cos \theta + qr \cos \theta + bd.$$

We conclude that  $a + c = b + d$  is equivalent to  $cpq + ars = bps + dqr$ .

Hence  $ABCD$  has an in circle if and only if

$$\frac{1}{h_1} + \frac{1}{h_3} = \frac{1}{h_2} + \frac{1}{h_4}.$$

6. From a set of 11 square integers, show that one can choose 6 numbers  $a^2, b^2, c^2, d^2, e^2, f^2$  such that

$$a^2 + b^2 + c^2 \equiv d^2 + e^2 + f^2 \pmod{12}.$$

**Solution:** The first observation is that we can find 5 pairs of squares such that the two numbers in a pair have the same parity. We can see this as follows:

Odd numbers	Even numbers	Odd pairs	Even pairs	Total pairs
0	11	0	5	5
1	10	0	5	5
2	9	1	4	5
3	8	1	4	5
4	7	2	3	5
5	6	2	3	5
6	5	3	2	5
7	4	3	2	5
8	3	4	1	5
9	2	4	1	5
10	1	5	0	5
11	0	5	0	5

Let us take such 5 pairs: say  $(x_1^2, y_1^2), (x_2^2, y_2^2), \dots, (x_5^2, y_5^2)$ . Then  $x_j^2 - y_j^2$  is divisible by 4 for  $1 \leq j \leq 5$ . Let  $r_j$  be the remainder when  $x_j^2 - y_j^2$  is divisible by 3,  $1 \leq j \leq 5$ . We have 5 remainders  $r_1, r_2, r_3, r_4, r_5$ . But these can be 0, 1 or 2. Hence either one of the remainders occur 3 times or each of the remainders occur once. If, for example  $r_1 = r_2 = r_3$ , then 3 divides  $r_1 + r_2 + r_3$ ; if  $r_1 = 0, r_2 = 1$  and  $r_3 = 2$ , then again 3 divides  $r_1 + r_2 + r_3$ . Thus we can always find three remainders whose sum is divisible by 3. This means we can find 3 pairs, say,  $(x_1^2, y_1^2), (x_2^2, y_2^2), (x_3^2, y_3^2)$  such that 3 divides  $(x_1^2 - y_1^2) + (x_2^2 - y_2^2) + (x_3^2 - y_3^2)$ . Since each difference is divisible by 4, we conclude that we can find 6 numbers  $a^2, b^2, c^2, d^2, e^2, f^2$  such that

$$a^2 + b^2 + c^2 \equiv d^2 + e^2 + f^2 \pmod{12}.$$