

Entrance Examination for CMI BSc (Mathematics & Computer Science) May 2011

Attempt all problems from parts A and C. Attempt any 7 problems from part B.

**Part A.** Choose the correct option and explain your reasoning briefly. Each problem is worth 3 points.

1. The word MATHEMATICS consists of 11 letters. The number of distinct ways to rearrange these letters is

- (A)  $11!$  (B)  $\frac{11!}{3}$  (C)  $\frac{11!}{6}$  (D)  $\frac{11!}{8}$

2. In a rectangle ABCD, the length BC is twice the width AB. Pick a point P on side BC such that the lengths of AP and BC are equal. The measure of angle CPD is

- (A)  $75^\circ$  (B)  $60^\circ$  (C)  $45^\circ$  (D) none of the above

3. The number of  $\theta$  with  $0 \leq \theta < 2\pi$  such that  $4 \sin(3\theta + 2) = 1$  is

- (A) 2 (B) 3 (C) 6 (D) none of the above

4. Given positive real numbers  $a_1, a_2, \dots, a_{2011}$  whose product  $a_1 a_2 \cdots a_{2011}$  is 1, what can you say about their sum  $S = a_1 + a_2 + \cdots + a_{2011}$  ?

- (A)  $S$  can be any positive number.  
(B)  $1 \leq S \leq 2011$ .  
(C)  $2011 \leq S$  and  $S$  is unbounded above.  
(D)  $2011 \leq S$  and  $S$  is bounded above.

5. A function  $f$  is defined by  $f(x) = e^x$  if  $x < 1$  and  $f(x) = \log_e(x) + ax^2 + bx$  if  $x \geq 1$ . Here  $a$  and  $b$  are unknown real numbers. Can  $f$  be differentiable at  $x = 1$ ?

- (A)  $f$  is not differentiable at  $x = 1$  for any  $a$  and  $b$ .  
(B) There exist unique numbers  $a$  and  $b$  for which  $f$  is differentiable at  $x = 1$ .  
(C)  $f$  is differentiable at  $x = 1$  whenever  $a + b = e$ .  
(D)  $f$  is differentiable at  $x = 1$  regardless of the values of  $a$  and  $b$ .

6. The equation  $x^2 + bx + c = 0$  has *nonzero* real coefficients satisfying  $b^2 > 4c$ . Moreover, exactly one of  $b$  and  $c$  is irrational. Consider the solutions  $p$  and  $q$  of this equation.

- (A) Both  $p$  and  $q$  must be rational.  
(B) Both  $p$  and  $q$  must be irrational.  
(C) One of  $p$  and  $q$  is rational and the other irrational.  
(D) We cannot conclude anything about rationality of  $p$  and  $q$  unless we know  $b$  and  $c$ .

7. When does the polynomial  $1 + x + \cdots + x^n$  have  $x - a$  as a factor? Here  $n$  is a positive integer greater than 1000 and  $a$  is a *real* number.

- (A) if and only if  $a = -1$   
(B) if and only if  $a = -1$  and  $n$  is odd  
(C) if and only if  $a = -1$  and  $n$  is even  
(D) We cannot decide unless  $n$  is known.

**Part B.** Attempt any 7 problems. Explain your reasoning. Each problem is worth 7 points.

1. In a business meeting, each person shakes hands with each other person, with the exception of Mr. L. Since Mr. L arrives after some people have left, he shakes hands only with those present. If the total number of handshakes is exactly 100, how many people left the meeting before Mr. L arrived? (Nobody shakes hands with the same person more than once.)

2. Show that the power of  $x$  with the largest coefficient in the polynomial  $(1 + \frac{2x}{3})^{20}$  is 8, i.e., if we write the given polynomial as  $\sum_i a_i x^i$  then the largest coefficient  $a_i$  is  $a_8$ .

3. Show that there are infinitely many perfect squares that can be written as a sum of six consecutive natural numbers. Find the smallest such square.

4. Let S be the set of all 5-digit numbers that contain the digits 1,3,5,7 and 9 exactly once (in usual base 10 representation). Show that the sum of all elements of S is divisible by 11111. Find this sum.

5. It is given that the complex number  $i - 3$  is a root of the polynomial  $3x^4 + 10x^3 + Ax^2 + Bx - 30$ , where  $A$  and  $B$  are unknown real numbers. Find the other roots.

6. Show that there is no solid figure with exactly 11 faces such that each face is a polygon having an odd number of sides.

7. To find the volume of a cave, we fit X, Y and Z axes such that the base of the cave is in the XY-plane and the vertical direction is parallel to the Z-axis. The base is the region in the XY-plane bounded by the parabola  $y^2 = 1 - x$  and the Y-axis. Each cross-section of the cave perpendicular to the X-axis is a square.

(a) Show how to write a definite integral that will calculate the volume of this cave.

(b) Evaluate this definite integral. Is it possible to evaluate it without using a formula for indefinite integrals?

8.  $f(x) = x^3 + x^2 + cx + d$ , where  $c$  and  $d$  are real numbers. Prove that if  $c > \frac{1}{3}$ , then  $f$  has exactly one real root.


9. A real-valued function  $f$  defined on a closed interval  $[a, b]$  has the properties that  $f(a) = f(b) = 0$  and  $f(x) = f'(x) + f''(x)$  for all  $x$  in  $[a, b]$ . Show that  $f(x) = 0$  for all  $x$  in  $[a, b]$ .

**Part C.** Explain your reasoning. Each problem is worth 10 points.

1. Show that there are exactly 16 pairs of integers  $(x, y)$  such that  $11x + 8y + 17 = xy$ . You need not list the solutions.

2. A function  $g$  from a set  $X$  to itself satisfies  $g^m = g^n$  for positive integers  $m$  and  $n$  with  $m > n$ . Here  $g^n$  stands for  $g \circ g \circ \cdots \circ g$  ( $n$  times). Show that  $g$  is one-to-one if and only if  $g$  is onto. (Some of you may have seen the term “one-one function” instead of “one-to-one function”. Both mean the same.)

3. In a quadrilateral  $ABCD$ , angles at vertices  $B$  and  $D$  are right angles.  $AM$  and  $CN$  are respectively altitudes of the triangles  $ABD$  and  $CBD$ . See the figure below. Show that  $BN = DM$ .



In this figure the angles  $ABC$ ,  $ADC$ ,  $AMD$  and  $CNB$  are right angles.