

# CHENNAI MATHEMATICAL INSTITUTE

Ph.D. Programme in Physics

Entrance Examination, 15 May 2013

[I] Consider a simple pendulum consisting of a bob of mass  $m$  suspended from a fixed support via a massless rigid rod of length  $l$ . It is free to oscillate (not necessarily through small angles) in a plane, under the influence of the Earth's constant acceleration due to gravity  $g$ .

1. Write the Lagrangian for this pendulum and plot the potential energy  $V(\theta)$ . Denote the angle made by the pendulum relative to its stable equilibrium position by  $\theta$  anti-clockwise. Choose the additive constant so that  $V(0) = -mgl$ .
2. Find Lagrange's equation for the pendulum. Obtain all static (time-independent) solutions and comment on their physical meaning and stability.
3. Find the momentum  $p_\theta$  conjugate to  $\theta$ . Starting from the Lagrangian, obtain the Hamiltonian expressed in terms of phase space variables via a Legendre transform.
4. What is the phase space of this system?
5. For energy  $-mgl \leq E < mgl$ , find the maximum angle of deflection  $\theta_0$ , and show that the time period of oscillation is as below. What is the origin of the pre-factor 4?

$$T = 4 \int_0^{\theta_0} \frac{d\theta}{\sqrt{\frac{2}{ml^2}[E + mgl \cos \theta]}}$$

6. Find  $\theta_0$  and  $T$  as  $E$  approaches  $mgl$  from below. What does the answer mean physically?

[II] Consider the wave function of a spin-two system given by

$$|\psi\rangle = \frac{a|S=2, S_z=2\rangle + b|S=2, S_z=1\rangle}{(|a|^2 + |b|^2)^{1/2}}$$

where

$$S_z|S=2, S_z=m\rangle = \hbar m|S=2, S_z=m\rangle$$

and

$$S^2|J=2, S_z=m\rangle = 6\hbar^2|S=2, S_z=m\rangle$$

and  $|S=2, S_z=m\rangle$  is normalised to unity. Let the Hamiltonian of the system be

$$H = \lambda \vec{S}^2 + \mu S_x^2$$

where  $\vec{S}$  is the spin operator and  $S_x$  is the  $x$ -component. In the above equations,  $a$  and  $b$  are complex constants and  $\lambda$  and  $\mu$  are real constants.

1. Suppose a measurement of  $S_z$  is made on  $|\psi\rangle$ . What are the possible outcomes and what is the probability of each?
2. What are the dimensions of  $\lambda$  and  $\mu$ ?
3. What are the conserved quantities of the Hamiltonian?
4. Find the expectation value of the Hamiltonian in the state  $|\psi\rangle$ .

[III]

1.  $A$  is a  $2 \times 2$  matrix

$$A = \begin{pmatrix} 1 & 2 \\ 0 & 3 \end{pmatrix}$$

Diagonalise  $A$  and find  $\exp(A)$ .

2. Show that the Fourier Transform of  $f(x) = \frac{1}{\mathbf{r}^2 + \lambda^2}$ , where  $\mathbf{r} = (x, y, z)$ ,  $\mathbf{r}^2 = x^2 + y^2 + z^2$  and  $\lambda^2 > 0$  is given by  $\tilde{F}(\mathbf{k}) = \frac{4\pi^2}{k} e^{-\lambda k}$ , where  $k = |\mathbf{k}|$ .

[IV] Consider a point particle of mass  $m$  and charge  $q$ .

1. Using the Lorentz force equation, find the transformation properties of electric field  $\vec{E}(\vec{r}, t)$ , magnetic field  $\vec{B}(\vec{r}, t)$ , vector potential  $\vec{A}(\vec{r}, t)$  and scalar potential  $\phi(\vec{r}, t)$ , under parity and (separately) time-reversal transformations.
2. Solve completely the force equation for the particle moving in a constant magnetic field in  $z$ -direction, giving the radius of the helix and the pitch angle.

[V] An ideal gas is at a temperature  $T_1$  and volume  $V_1$ . The gas is taken through an isobaric (constant pressure) process to a state of higher temperature  $T_2$ . It is then taken via an isochoric (constant volume) process to a state of temperature  $T_1$ , and finally back to the initial state in an isothermal (constant temperature) process.

1. Calculate the amount of heat transferred ( $Q$ ) to the gas in the cycle.
2. Same as above, but in the reverse cycle.
3. What would be the result if  $\delta Q$  were an exact differential?
4. Calculate the work done by the gas during the cycle. Is it equal to  $Q$ ? Why?
5. Draw a P-V diagram to illustrate the cycle.
6. Comment on the change in entropy, if any, during the cycle.