

1. The maximum value of $|z|$ when z satisfies the condition $|z + \frac{1}{z}| = 4$ is
 A) $2 - \sqrt{5}$ B) $2 + \sqrt{5}$ C) $4 - \sqrt{5}$ D) $4 + \sqrt{5}$
2. If $1, \omega_1, \omega_2, \dots, \omega_9$ are the 10th roots of unity, then $(1 + \omega_1)(1 + \omega_2) \cdots (1 + \omega_9)$ is
 A) 0 B) 1 C) -1 D) 9
3. If x is a real number, then $(x - 1)^2 + (x - 2)^2 + \cdots + (x - 100)^2$ is least when x is
 A) 50 B) 100 C) 101 D) $\frac{101}{2}$
4. The sum $100C_0 + 101C_1 + 102C_2 + \cdots + 150C_{50}$ is
 A) $200C_{100}$ B) $201C_{50}$ C) $201C_{100}$ D) $151C_{50}$
5. If $A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ a & b & -1 \end{pmatrix}$ then A^{101} is
 A) I B) $A - I$ C) A D) $(a + b)(A - I)$
6. The value of the determinant $\begin{vmatrix} 1 & \log_5 10 & \log_5 15 \\ \log_{10} 5 & 1 & \log_{10} 15 \\ \log_{15} 5 & \log_{15} 10 & 1 \end{vmatrix}$ is
 A) 0 B) 1
 C) $\log_5 150 + \log_{10} 75 + \log_{15} 50$ D) $\log_5 25 + \log_{10} 20 + \log_{15} 15$
7. For what value of λ will the equation $\lambda x^2 - 10xy + 12y^2 + 5x - 16y - 3 = 0$ represent a pair of straight lines
 A) 4 B) 2 C) -2 D) 3
8. The equation of a tangent to the circle $x^2 + y^2 - 2x - 6y - 12 = 0$ is
 A) $\sqrt{3}(x - 2) + (y - 3) = 0$
 B) $\sqrt{3}(x - 2) + (y - 3) = 5$
 C) $\sqrt{3}(x - 2) + (y - 3) = 10$
 D) $(x - 2) + \sqrt{3}(y - 3) = 5$

9. The director circle of the ellipse $\frac{x^2}{16} + \frac{y^2}{9} = 1$ is
- A) $x^2 + y^2 = 16$ B) $x^2 + y^2 = 9$
 C) $x^2 + y^2 = 7$ D) $x^2 + y^2 = 25$
10. The angle between the planes $2x - y + z = 6$ and $x + y + 2z = 3$ is
- A) π B) $\frac{\pi}{2}$ C) $\frac{\pi}{3}$ D) $\frac{\pi}{6}$
11. The equation of the perpendicular bisector of the straight line joining the points $(2, 3)$ and $(1, 2)$ is
- A) $x - y + 4 = 0$ B) $x - y - 2 = 0$
 C) $x + y - 4 = 0$ D) $x + y - 2 = 0$
12. The spheres $x^2 + y^2 + z^2 = 25$ and $x^2 + y^2 + z^2 - 24x - 40y - 18z + 225 = 0$
- A) touch internally
 B) touch externally
 C) do not touch each other
 D) intersect each other
13. $\cos 2x + a \sin x = 2a - 7$ possesses a solution for
- A) all a B) $a > 6$ C) $a < 2$ D) $a \in [2, 6]$
14. The lowest degree of the polynomial with real coefficients having roots $2, -3, 2 + i, 1 + i$ is
- A) 2 B) 4 C) 6 D) 8
15. Let $f(x) = 6x + 5$. If f_n denotes the function $f \circ f \circ \dots \circ f$ n times then $f_{15}(5)$ is
- A) $6^{15} - 1$ B) $6^{15} + 1$ C) $6^{16} - 1$ D) $5(6^{15} + 1)$
16. If $f(x) = 2^x + 2^{x+1} + \dots + 2^{x+9}$ then $f'(2)$ is
- A) $1023 \log_e 16$ B) $1023 \log_e 8$ C) $1023 \log_e 4$ D) $1023 \log_e 2$

17. If $f(x) = \min\{x, x^2\}$ for every real value of x , then which one of the following is not true

- A) f is continuous for all x
- B) f is differentiable for all x
- C) $f'(x) = 1$ for all $x > 1$
- D) one of the above statement is wrong

18. If $\int_0^{\frac{\pi}{2}} \cos^n x dx = A$, then the value of $n \int_{\frac{\pi}{2}}^0 \sin^n x dx$ is

- A) $-A$
- B) A
- C) nA
- D) $-nA$

19. If $\int_0^x f(t) dt = x + \int_x^1 tf(t) dt$ then the value of $f(1)$ is

- A) $\frac{1}{2}$
- B) $-\frac{1}{2}$
- C) 1
- D) -1

20. The general solution of the equation $(e^{-x} + \sin y)dx + \cos y dy = 0$ is

- A) $x + e^{-x} \cos y + C = 0$
- B) $x - e^{-x} \sin y + C = 0$
- C) $x + e^x \sin y + C = 0$
- D) $x - e^x \sin y + C = 0$

21. $\lim_{n \rightarrow \infty} \{\sqrt{n^2 + n} - n\}$ is

- A) 0
- B) 1
- C) $\frac{1}{2}$
- D) ∞

22. $\lim_{n \rightarrow \infty} (n^{\frac{1}{n}} - 1)^n$ is

- A) 1
- B) 0
- C) e
- D) ∞

23. Which of the following series is divergent

- A) $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$
- B) $\sum_{n=1}^{\infty} \frac{1}{n \log(n+1)}$
- C) $\sum_{n=1}^{\infty} \frac{\sqrt{n+1} - \sqrt{n}}{n}$
- D) $\sum_{n=1}^{\infty} \frac{1}{n^2}$

24. Which of the following sequence is convergent for all x in $[0, 1]$, but is not uniformly convergent on $[0, 1]$?

- A) $\{\frac{\sin nx}{\sqrt{n}}\}$ B) $\{\sin nx\}$ C) $\{x^n(1+x)^{-n}\}$ D) $\{x^n\}$

25. If $A = \lim_{x \rightarrow 0} x \sin \frac{1}{x}$ and $B = \lim_{x \rightarrow \infty} x \sin \frac{1}{x}$, then

- A) $A = B = 0$ B) $A = 0$ and $B = \infty$
 C) $A = 0$ and $B = 1$ D) $A = 1$ and $B = \infty$

26. Let $[x]$ denote the greatest integer not exceeding x , then the value of the Riemann Stielgies integral $\int_0^2 x^2 d[x]$ is equal to

- A) 1 B) 3 C) 5 D) 0

27. Let the function f be defined on \mathbb{R} by

$$f(x) = \begin{cases} 0, & \text{if } x \text{ is rational} \\ x, & \text{Otherwise} \end{cases}$$

Let μ be the Lebesgue measure on $[0, 1]$, then the Lebesgue integral $\int_0^1 f d\mu$ has the value

- A) 1 B) 0 C) $\frac{1}{2}$ D) 2

28. Let $f(x) = \begin{cases} 1, & \text{if } x \text{ is rational} \\ -1, & \text{if } x \text{ is irrational} \end{cases}$

Then which of the following function is Riemann integrable on $[0, 1]$

- A) f B) $|f|$ C) f^+ D) f^-

29. If the radius of convergence of the power series $\sum_{n=0}^{\infty} a_n z^n$ is R , then the radius of convergence of the power series $\sum_{n=0}^{\infty} n^2 a_n z^n$ is

- A) R B) $2R$ C) $\frac{R}{2}$ D) R^2

30. Which of the following power series represent the principal branch of $\log(1+z)$?
- A) $z - \frac{z^2}{2} + \frac{z^3}{3} - \dots$ B) $z + \frac{z^2}{2} + \frac{z^3}{3} + \dots$
 C) $1 + z + \frac{z^2}{2} + \dots$ D) $1 - z + \frac{z^2}{2} - \dots$
31. Let γ be the path defined by $\gamma(t) = e^{4\pi it}$, $0 \leq t \leq 1$. Then the value of the integral $\int_{\gamma} \frac{dz}{z}$ is
- A) $2\pi i$ B) $4\pi i$ C) 0 D) $-2\pi i$
32. The singularity of the function $\frac{1 - \cos z}{z^2}$ at $z = 0$ is
- A) a simple pole B) a pole of order 2
 C) a removable singularity D) an essential singularity
33. Let γ be a positively oriented unit circle, then $\int_{\gamma} \frac{\sin z}{z^2} dz$ has the value
- A) $2\pi i$ B) 0 C) $-2\pi i$ D) $4\pi i$
34. At $z = 0$, the function $f(z) = \frac{1}{z} + \frac{1}{z^2} + e^{\frac{1}{z}}$ has
- A) an essential singularity B) a simple pole
 C) a pole of order 2 D) a removable singularity
35. The radius of convergence of the power series $\sum_{n=0}^{\infty} \frac{n^2 z^{2n}}{2^n}$ is
- A) $\frac{1}{\sqrt{2}}$ B) 2 C) $\sqrt{2}$ D) $\frac{1}{2}$
36. Which of the following subsets of the complex plane is simply connected?
- A) $\{z : |z| > 1\}$
 B) $\{z : |z - 1| \leq 2\} \cup \{z : |z + 1| \leq 2\}$
 C) $\{z : 0 < |z| < 1\}$
 D) $\{z : |z - 1| > 1\}$

37. Let T be the Möbius transformation defined by $T(z) = \frac{z+i}{iz+1}$. Then T maps the real axis $\{z : \operatorname{Im} z = 0\}$ onto
- A) the imaginary axis $\{z : \operatorname{Re} z = 0\}$
 - B) the unit circle $\{z : |z| = 1\}$
 - C) the line $\{z : \operatorname{Re} z = 1\}$
 - D) the circle $\{z : |z - i| = 1\}$
38. Let $f(z) = \sin \frac{\pi}{z}$, $z \in \mathbb{C}$, $z \neq 0$. Then which of the following statements is incorrect.
- A) $f(z)$ has infinite number of zeros in \mathbb{C}
 - B) $z = 0$ is an essential singularity of f
 - C) $\lim_{|z| \rightarrow \infty} f(z) = 0$
 - D) $f(z)$ is bounded in the annulus $\{z : 0 < |z| < 1\}$
39. The residue at $z = 1$ of the function $\frac{1}{(z-1)(z-3)^2}$ is
- A) 2
 - B) 0
 - C) $\frac{1}{4}$
 - D) 4
40. The coefficient of $\frac{1}{z}$ in the Laurent series expansion of $f(z) = \frac{1}{z(z-1)}$ in the region $1 < |z| < \infty$ is
- A) 1
 - B) 0
 - C) -1
 - D) 2
41. Which of the following permutations is even
- A) $\begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 4 & 5 & 1 \end{pmatrix}$
 - B) $\begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 5 & 3 & 4 & 2 & 1 \end{pmatrix}$
 - C) $\begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 1 & 4 & 5 & 3 \end{pmatrix}$
 - D) $\begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 1 & 2 & 5 & 4 \end{pmatrix}$
42. If $a + bi$ with $a, b \in \mathbb{Z}$ is a unit in the ring $\mathbb{Z}[i]$ of Gaussian integers, then which of the following is true
- A) $a = 1$
 - B) $a = -1$
 - C) $b = 1$
 - D) $ab = 0$
43. Which of the following groups is cyclic
- A) $\mathbb{Z}_6 \oplus \mathbb{Z}_8$
 - B) $\mathbb{Z}_3 \oplus \mathbb{Z}_{16}$
 - C) $\mathbb{Z}_4 \oplus \mathbb{Z}_{12}$
 - D) $\mathbb{Z}_2 \oplus \mathbb{Z}_{24}$

44. The order of the element $(2, 2)$ in the group $\mathbb{Z}_4 \oplus \mathbb{Z}_6$ is
 A) 2 B) 4 C) 6 D) 8
45. For which of the following numbers all groups of that order are abelian
 A) 6 B) 8 C) 12 D) 25
46. Which of the following pair of groups are isomorphic
 A) \mathbb{Z}_{24} and $\mathbb{Z}_8 \oplus \mathbb{Z}_3$ B) \mathbb{Z}_{25} and $\mathbb{Z}_5 \oplus \mathbb{Z}_5$
 C) \mathbb{Z}_4 and $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ D) \mathbb{Z}_{20} and $\mathbb{Z}_2 \oplus \mathbb{Z}_{10}$
47. Which of the following maps is a homomorphism on the ring $\mathbb{Z} \times \mathbb{Z}$
 A) $\phi(x, y) = (2x, 2y)$ B) $\phi(x, y) = (x + y, 0)$
 C) $\phi(x, y) = (2x, 3y)$ D) $\phi(x, y) = (y, x)$
48. Which of the following is a unit in the ring $\mathbb{Z}(\sqrt{2}) = \{a + b\sqrt{2} : a, b \in \mathbb{Z}\}$
 A) $3 + 2\sqrt{2}$ B) $2 + 3\sqrt{2}$ C) $2 + \sqrt{2}$ D) $1 + 2\sqrt{2}$
49. Which of the following equations has a solution in \mathbb{Z}_{18}
 A) $3x = 5$ B) $4x = 3$ C) $5x = 4$ D) $6x = 7$
50. Which of the following polynomials is not irreducible in $\mathbb{Z}_3[x]$
 A) $x^2 + 1$ B) $x^2 + x + 2$
 C) $x^3 + x^2 + 2$ D) $x^3 + x + 1$
51. Which of the following is an ideal in the ring $F[x]$ of all polynomials over a field F
 A) set of all polynomials in $F[x]$ of degree > 1
 B) set of all polynomials in $F[x]$ of degree ≤ 1
 C) set of all polynomials in $F[x]$ without constant term
 D) set of all polynomials $f(x) \in F[x]$ such that $f(0) \neq 0$
52. The degree of the field extension $[\mathbb{Q}(\sqrt{2} + \sqrt{3}), \mathbb{Q}]$ is
 A) 1 B) 2 C) 3 D) 4

53. Which of the following statement is not true about an algebraically closed field K
- A) Every non constant polynomial in $K[x]$ has a zero in K
 B) Every polynomial in $K[x]$ of degree n has a factorization into n linear factors in $K[x]$
 C) Irreducible polynomials in $K[x]$ have degree ≤ 1
 D) Every extension of K is an algebraic extension
54. Let $K = \mathbb{Q}(\alpha)$ where α is the real cube root of 2, then the order of the automorphism group $\text{Aut}(K, \mathbb{Q})$ is
- A) 1 B) 2 C) 4 D) 6
55. Let σ be an automorphism in $\text{Aut}(\mathbb{Q}(\sqrt{2}, \sqrt{3}) : \mathbb{Q})$. Then which of the following can not hold
- A) $\sigma(\sqrt{2}) = -\sqrt{2}$ B) $\sigma(\sqrt{2}) = \sqrt{3}$
 C) $\sigma(\sqrt{2} + \sqrt{3}) = \sqrt{2} - \sqrt{3}$ D) $\sigma(\sqrt{2} + \sqrt{3}) = -\sqrt{2} + \sqrt{3}$
56. In the vector space \mathbb{R}^3 over \mathbb{R} , W is the subspace given by $W = \{(x_1, x_2, x_3) : x_1 + x_2 + x_3 = 0\}$. Then $\dim W$ is
- A) 0 B) 1 C) 2 D) 3
57. Which of the following is a linearly independent set in \mathbb{R}^2
- A) $\{(1, -1), (-2, 2)\}$ B) $\{(1, -1), (3, -1)\}$
 C) $\{(1, 2), (2, 4)\}$ D) $\{(3, 1), (-3, -1)\}$
58. Which of the following is an eigen vector of the matrix $A = \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix}$
- A) $\begin{bmatrix} 1 \\ 2 \end{bmatrix}$ B) $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$ C) $\begin{bmatrix} 3 \\ 0 \end{bmatrix}$ D) $\begin{bmatrix} 0 \\ 2 \end{bmatrix}$
59. Which of the following matrix is diagonalizable
- A) $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$ B) $\begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$ C) $\begin{bmatrix} 2 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 3 \end{bmatrix}$ D) $\begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$
60. Let T from \mathbb{R}^2 to \mathbb{R}^3 be defined by $T(x, y) = (x + y, x + y, 0)$. Then rank T is
- A) 0 B) 1 C) 2 D) 3

61. With usual metric in \mathbb{R} which of the following subspaces of \mathbb{R} is complete
- A) the rationals in \mathbb{R}
 - B) the irrationals in \mathbb{R}
 - C) the closed interval $[0, 1]$
 - D) the open interval $(0, 1)$
62. With usual topology on the spaces concerned which of the following spaces is not connected?
- A) $\{z \in \mathbb{C} : |z| < 1\}$
 - B) $\{x \in \mathbb{R} : |x| < 1\}$
 - C) $\{z \in \mathbb{C} : |z| > 1\}$
 - D) $\{x \in \mathbb{R} : |x| > 1\}$
63. Which of the following is not a property of \mathbb{R} (with usual topology)
- A) second countability
 - B) compactness
 - C) separability
 - D) local compactness
64. Which among the following topologies on \mathbb{R} is an example of a topology not induced by a pseudo metric?
- A) usual topology
 - B) discrete topology
 - C) indiscrete topology
 - D) cofinite topology
65. Which of the following functions $d : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is not a metric
- A) $d(x, y) = |x - y|$
 - B) $d(x, y) = 2|x - y|$
 - C) $d(x, y) = \frac{|x - y|}{1 + |x - y|}$
 - D) $d(x, y) = |x - y|^2$
66. Let X be a topological space and let A, B be subsets of X . Then it is not always true that
- A) $\overline{\bar{A}} = \bar{A}$
 - B) $\overline{(A \cup B)} = \bar{A} \cup \bar{B}$
 - C) $\overline{(A \cap B)} = \bar{A} \cap \bar{B}$
 - D) $\bar{X} = X$
67. With the usual topology, which of the following subspaces of \mathbb{R} is not homeomorphic to $(0, 1)$?
- A) $\{x | x > 0\}$
 - B) $[0, 1]$
 - C) \mathbb{R}
 - D) $(-1, 1)$

68. Let X be a metric space. Three of the following properties of X are equivalent to each other, pick the odd one out
- A) X is compact
 - B) X is sequentially compact
 - C) X has the Bolzano-Weierstrass property
 - D) X is totally bounded
69. Let \mathbb{R} be the space of real numbers with usual topology. Which of the following subspaces of \mathbb{R} is compact?
- A) $(0, 1)$
 - B) $[0, 1] \cup [2, 3]$
 - C) $[0, 1)$
 - D) set of all rationals in \mathbb{R}
70. Let (X, τ) be the Sierpinski topology with $X = \{a, b\}$, $\tau = \{\phi, \{a\}, X\}$. Then X is not a
- A) compact space
 - B) connected space
 - C) T_0 space
 - D) T_1 space
71. Let X be the normed linear space of square summable real sequences with $\| \cdot \|_2$ and Y be the subspace generated by the elements $(1, 0, 0, \dots)$ and $(0, 1, 0, \dots)$. If $U = \{x \in X : \|x\|_2 < 1\}$ Then
- A) $Y + U$ is open in X
 - B) $Y + U$ is closed in X
 - C) $Y + U$ is neither open nor closed in X
 - D) $Y + U$ is not bounded in X
72. Let X be the complex normed linear space of summable sequences of complex numbers with norm $\| \cdot \|_1$ and $Y = \{x \in X : \|x\|_1 \leq 1\}$ then
- A) Y is compact and convex
 - B) Y is compact but not convex
 - C) Y is neither compact nor convex
 - D) Y is convex but not compact
73. Let $X = C_{00}$, the space of all real sequences which have only finitely many nonzero members, and f be the linear functional on X defined by $f(x(1), x(2), \dots) = x(1) + x(2) + \dots$ for $x = (x(1), x(2), \dots) \in X$. Then f is continuous
- A) with respect to $\| \cdot \|_1$ and $\| \cdot \|_2$ but not with respect to $\| \cdot \|_\infty$
 - B) with respect to $\| \cdot \|_1$ and $\| \cdot \|_\infty$ but not with respect to $\| \cdot \|_2$
 - C) with respect to $\| \cdot \|_2$ and $\| \cdot \|_\infty$ but not with respect to $\| \cdot \|_1$
 - D) with respect to $\| \cdot \|_1, \| \cdot \|_2$ and $\| \cdot \|_\infty$

74. Let $X = C_{00}$ with $\|\cdot\|_{\infty}$ and $F : X \rightarrow l^{\infty}$ be a bounded linear map. Then there is a bounded linear map $G : C_0 \rightarrow l^{\infty}$ such that

- A) G is unique, $G/C_{00} = F$ and $\|F\| < \|G\|$
- B) G is unique, $G/C_{00} = F$ and $\|F\| = \|G\|$
- C) $G/C_{00} = F$ and $\|F\| = \|G\|$ but G is not necessarily unique
- D) G is unique, $R(G) = R(F)$ and $\|F\| < \|G\|$

75. Let X be a normed linear space and Y be a subspace of X with basis $\{y_1, y_2, \dots, y_n\}$. Let x'_1, x'_2, \dots, x'_n be linear functionals with

$$x'_i(y_j) = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

If $Z = \{x : x'_j(x) = 0, \text{ for } j = 1, 2, \dots, n\}$ then which one of the following is not correct?

- A) $Y \cap Z = \{0\}$
- B) $Y + Z = X$
- C) Z is open
- D) Z is closed

76. If H is the Hilbert space of square summable sequences of complex numbers and if $x = (x(1), x(2), \dots) \in H$ has the property that $2 \sum_{i=1, i \neq j}^{\infty} |x(i)|^2 + |x(j) - 1|^2 + |x(j) + 1|^2 = 18$ then $\|x\|$ is equal to

- A) 1
- B) 2
- C) $2\sqrt{2}$
- D) 4

77. Let H be the complex Hilbert space of square summable sequences of complex numbers and $T : H \rightarrow H$ be defined $T(x(1), x(2), \dots) = (0, x(1), x(2), \dots)$ for $x = (x(1), x(2), \dots) \in H$. Then which one of the following is not correct?

- A) T is bounded
- B) $\|T\| = 1$
- C) T is one-one but not onto
- D) T is one-one and onto

78. Let M be a closed subspace of a complex Hilbert space H . Let P and Q be orthogonal projections of H onto M and M^{\perp} respectively. Then the set of all values of α, β such that $\alpha P + \beta Q$ is selfadjoint is

- A) ϕ
- B) $\{1\}$
- C) the set of all real numbers
- D) set of all complex numbers

79. Let H be the real Hilbert space $L^2([0, 2\pi])$ and f be a linear functional on H defined by $f(x) = \int_0^{2\pi} x \sin 2x dx$. Then $\|f\|$ is
- A) 1 B) π C) 2π D) $\sqrt{\pi}$
80. Let X_1 and X_2 be closed subspaces of a Hilbert space H and let P_1 and P_2 be orthogonal projections on X_1 and X_2 respectively. If $\langle x, y \rangle = 0$ for all $x \in X_1, y \in X_2$ then which one of the following is not correct?
- A) $X_1 + X_2$ is a closed subspace of H
B) $P_1 - P_2$ is an orthogonal projection
C) $(P_1 - P_2)^2$ is an orthogonal projection
D) $P_1 + P_2$ is an orthogonal projection
-