



iQuest Scholastic & Mental Aptitude Reward Test

FOR CLASS 12TH (EXCEL) SAMPLE TEST (MED.)

Time: 1 Hr	Max Marks: 240

SYLLABUS & SCHEME						
SUBJECTS	Qs.	SYLLABUS				
PHYSICS	15	Units & Dimensions, Kinematics, NLM, WPE, Circular Motion,				
		Collision, COM, Rotation				
CHEMISTRY	15	Atomic Structure, Mole Concept, Gaseous State,				
		Periodic Classification, Chemical Bonding, Thermodynamics,				
		Chemical Equilibrium, Ionic Equilibrium, Redox Reactions				
BOTANY	15	Morphology, Anatomy, Classification, Plantae				
ZOOLOGY	15	Animal Kingdom, Biomolecules, Animal Morphology, Animal				
		Tissue, Digestion and Absorption, Breathing and Exchange of gases				


INSTRUCTIONS TO CANDIDATE

- Each subject in this paper consists of multiple choice questions with only one correct answer. +4 marks will be awarded for correct answer and -1 mark for wrong answer.
- Please read the instructions given for each question carefully and fill the correct answer against the question numbers on the answer sheet in the respective subject.
- Use blue or black ball point pen to darken the appropriate circle & mark should completely fill the circle.
- The Question paper contains blank spaces for your rough work. No additional sheet will be provided for rough work.
- Blank papers, Clipboards, Log Tables, Slide rule, Calculators, Cellular phones, Pagers and Electronic gadgets in any form are not allowed.
- Write your Name, Student ID in the block at the top of the Answer Sheet. Also write your Name & Student ID in the space provided on this cover page of question paper.
- This test paper is just an indicative of the actual test. Total number and type of questions in actual test may vary.




PHYSICS


- 1. A particle is thrown vertically upwards from the surface of the earth. Let T_p be the time taken by the particle to travel from a point P above the earth to its highest point and back to the point P. Similarly, let T_o be the time taken by the particle to travel from another point Q above the earth to its highest point and back to the same point Q. If the distance between the points P and Q is H, the expression for acceleration due to gravity in terms of T_P, T_O and H, is :-
 - (A) $\frac{6H}{T_P^2 + T_O^2}$
- (B) $\frac{8H}{T_p^2 T_0^2}$
- (C) $\frac{2H}{T_P^2 + T_0^2}$
- (D) $\frac{H}{T_{\rm p}^2 T_{\rm o}^2}$
- 2. An aeroplane is travelling horizontally at a height of 2000 m from the ground. The aeroplane, when at a point P, drops a bomb to hit a stationary target Q on the ground. In order that the bomb hits the target, What angle θ must the line PQ make with the vertical ? [$g = 10 \text{ ms}^{-2}$]



- (A) 15°
- (B) 30°
- (C) 90°
- (D) 45°
- **3.** For the following system




- (A) Acceleration of the system = $2m/s^2$
- (B) $T_1 = 20 \text{ N}$
- (C) $T_2 = 10N$
- (D) $T_2 > T_1$
- 4. For shown situation in figure (Assume: $g = 10 \text{ ms}^{-2}$)




- (A) The acceleration of the block A is 1m/s²
- (B) The acceleration of the block B is 3m/s²

- (C) The time taken for the front face of A lining up with the front face of B is 0.25 sec
- (D) The time taken for the front face of A lining up with the front face of B is 0.50 sec
- 5. Two equal masses are placed as shown in the figure. Friction at the pulley is negligible. If coefficient of sliding friction of the mass on the horizontal surface is 0.2 and if the hanging mass is just released from position of rest, the acceleration of the system, is (take $g = 9.8 \text{ m/s}^2$):-




- (A) 1 m/s^2
- (B) 2 m/s^2
- (C) 3.92 m/s^2
- (D) 4 m/s^2
- 6. A block of mass 5kg is placed on horizontal surface, and a pushing force 20N is acting on block as shown in fig. If coefficient of friction between block and surface is 0.2 then frictional force and speed of block after 15 sec, are respectively:-

(Given
$$g = 10 \text{ m/s}^2$$
)




- (A) $(5+2\sqrt{2})$ N, 3.25 ms⁻¹
- (B) $(10+2\sqrt{2})$ N, 3.25 ms⁻¹
- (C) $(5+2\sqrt{2})$ N, 3.94 ms⁻¹
- (D) $(10 + 2\sqrt{2})$ N, 3.94 ms⁻¹
- 7. A particle of mass 'M' falls from height 'h' and gets stick after collision, with identical particle lying on sand. After sticking, both particles moves a distance d in sand, then the work done against retarding force of sand is :-
 - (A) $\frac{Mgh}{2} + 2Mgd$ (B) $\frac{Mgh}{2} + Mgd$
 - (C) $\frac{Mgh}{2} 2Mgd$ (D) $\frac{Mgh}{2} Mgd$

- 8. A body is dropped from height 8m. After striking the surface it rises to 6m, the fractional loss in kinetic energy during impact, is (Assuming the frictional resistance to be negligible)
 - (A) $\frac{1}{2}$
- (C) $\frac{1}{5}$
- (D) $\frac{1}{7}$
- 9. A body of mass 0.8 kg has initial velocity $(3\hat{i} - 4\hat{j})$ m/sec. and final velocity $(-6\hat{j} + 2\hat{k})$ m/ sec, the change in kinetic energy of the body is:
 - (A) 2 J
- (B) 3 J
- (C) 4 J
- (D) 6 J
- A chain of mass m and length L is held on a fric-**10.** tionless table in such a way that its $\frac{1}{n}$ th part is hanging below the edge of table. The work done to pull the hanging part of chain is :-
 - (A) $\frac{mgL^2}{2n^2}$
- (B) zero
- (C) $\frac{mgL}{2n}$
- The angle between angular momentum and linear momentum for a particle in motion is :
 - o
- (B) 90°
- (C) 45°
- (D) 180°
- 12. Two identical rods each of mass M and length L are kept according to figure. The moment of inertia of rods about an axis passing through O and perpendicular to the plane of rods, is:-




- (A) $\frac{1}{3}ML^2$
- (B) $\frac{2}{3}ML^2$
- (C) $2ML^2$
- (D) $\frac{1}{2}ML^2$
- A flywheel rotates with a uniform angular acceleration. Its angular velocity increases from 20π rad/ s to 40π rad/s in 10 seconds. The number of rotations, it made in this period are:

- (A) 100
- (B) 150
- (C) 200
- (D) 250
- Three identical rings of mass 'M' and radius 'R' 14. are placed shown in figure. The moment of inertia about axis xx' is:



- (A) $\frac{5}{2}MR^2$
- (B) $\frac{7}{2}MR^2$
- (C) $\frac{3}{2}MR^2$ (D) $\frac{9}{2}MR^2$
- A disc is rotating with angular velocity (ω) about **15.** its axis (without any translation push) on a smooth surface: The directions and magnitudes of velocity at points B and A are:



- (A) $V_A = +\frac{\omega R}{2}$ (Towards right), $V_B = -\omega R$ (Towards left)
- (B) $V_A = -\omega R$ (Towards right), $V_B = \frac{\omega R}{2}$ (Towards left)
- (C) $V_A = +\frac{\omega R}{4}$ (Towards right), $V_B = -\omega R$ (Towards left)
- (D) $V_A = +\frac{\omega R}{2}$ (Towards right), $V_B = \frac{\omega R}{2}$ (Towards left)

CHEMISTRY

16. For the redox reaction,

$$\mathsf{MnO_4^-} + \mathsf{C}_2\mathsf{O}_4^{2-} + \mathsf{H}^+ \longrightarrow \mathsf{Mn}^{2+} + \mathsf{CO}_2 + \mathsf{H}_2\mathsf{O}$$

The correct coefficients of the reactants for the balanced reaction are -

- $MnO_4^ C_2O_4^{2-}$ H^+
- (A) 2 5 16
- (B) 16 5 2
- (C) 5 16 2
- (D) 2 16 5
- The oxidation state of S in Na₂S₄O₆ is -**17.**
 - (A) + 2
- (B) +4
- (C) +6
- (D) + 2.5
- 18. The empirical formula of a compound is CH. Its molecular weight is 78. The molecular formula of the compound will be:
 - (A) C₂H₂
- (B) C_3H_3
- (C) C_4H_4
- 19. One mole of a non-ideal gas undergoes a change of state (2.0 atm, 3.0L, 95K) \rightarrow (4.0 atm, 5.0L, 245 K) with a change in internal energy, U = 30.0 L-atm. The change in enthalpy (H) of the process in L-atm is:
 - (A) 40.0
- (B)42.3
- (C) 44.0
- (D) Not defined, because pressure is not constant
- Standard molar heats of formation of ethane 20. (C_2H_6) , $CO_2(g)$ and $H_2O(l)$ are -21.1, -94.1and -68.3 kcal respectively. The standard enthalpy of combustion of ethane is
 - (A) 372 kcal
- (B) -472 kcal
- (C) -372 kcal
- (D) -68.3 kcal
- 21. At infinite dilution, the percentage ionisation for both strong and weak electrolytes is
 - (A) 1%
- (B) 20%
- (C) 50%
- (D) 100%
- Which of the following pairs do not form peroxide
 - (A) Na,K
- (B) Li, Mg
- (C) Ca,Sr
- (D) Ca, Ba
- 23. Covalent compounds are generally in water
 - (A) Soluble
- (B) Insoluble
- (C) Dissociated
- (D) Hydrolysed
- $\frac{K_p}{K}$ for the given reaction will be
 - $2CO(g) + O_2(g)$ $2CO_2(g)$ is
 - (A) RT
- (B) $\frac{1}{PT}$

- (C) $\frac{1}{\sqrt{RT}}$
- (D) \sqrt{RT}
- 25. When equal volumes of pH = 4 and pH = 6 are mixed together then the pH of the resulting solution will be [Take $\log 5 = 0.7$]
 - (A) 5.75
- (B) 4.3

(C) 5

- (D) 5.3
- If the $k_{_{SD}}$ for AgCl is 1.8×10^{-10} , what would be its **26.** molar solubility in 1M AgNO₃
 - (A) 1.8×10^{-10}
- (B) 1.8×10^{-9}
- (C) 3.2×10^{-20}
- (D) 1.34×10^{-10}
- For the given equilibrium

$$Ag^+ + 2NH$$

$$Ag^+ + 2NH_3$$
 $Ag(NH_3)_2^+$ $k_1 = 1.8 \times 10^7$

$$Ag^+ + Cl$$

$$Ag^{+} + Cl^{-}$$
 $AgCl$ $k_{2} = 5.6 \times 10^{9}$

Than, what would be the equilibrium constant for process given below

$$AgCl + 2NH_{2}$$

$$[Ag(NH_3)_2]^+ + Cl^-$$

- (A) 10^{-17}
- (B) 3.1×10^{-22}
- (C) 3.2×10^{-3}
- (D) 10^{17}
- 28. If the radius of first orbit of H is r, then the radius of first Bohr's orbit of Li2+ will be -
 - (A) 3 r
- (B) 27 r
- (C) $\frac{r}{27}$
- (D) $\frac{r}{2}$
- For the non-zero volume of molecules having no forces of attraction, the variaion of compressibility

factor $Z = \frac{PV}{PT}$ with pressure is given by the graph

- (A) I
- (B) II
- (C) III
- (D) IV
- Pressure
- An exothermic reaction is represented by the graph **30.**

BOTANY

31.	Elaters help in dispersal of spores of (A) Riccia (B) Marcha (C) Dryopteris (D) Funario	ntia	to plant structure is co				
32.	Which of following type of anther Malvaceae? (A) Monothecous (B) Ditheco (C) Polythecous (D) Without	is found in	transpiration (B) Passage cells he cortex to phloem	ata but lenticels carry out lp in transfer of food from nts possess cytoplasm but no			
33.	Fascicular cambium is the cambium bundle of: (A) monocot stem (C) monocot leaf (D) dicot leaf	em of	nuclei (D) The shoot apica centre	l meristem has a quiescent			
34.	Companion cells in plants are associated (A) vessels (B) sperms (C) sieve elements (D) guard companion cells in plants are associated (B) sperms (D) guard companion cells in plants are associated (B) sperms (C) sieve elements (D) guard companion cells in plants are associated (C) sieve elements (D) guard companion cells in plants are associated (E) sperms (E) s	ated with:	lysis, are called: (A) lysozymes (C) lytic	teria multiply and cause their (B) lipolytic (D) lysogenic			
35.	Mosses and ferns are found in moi places because both: (A) require presence of water for fert (B) do not need sunlight for photosyn (C) depend for their nutrition on mic which can survive only at low temper	ilization thesis ro-organisms rature	Choose the correct securve for bacteria (A) lag, log, stationary (B) lag, log, stationary (C) stationary, lag, log	Choose the correct sequence of stages of growth			
36.	(D) cannot compete with sun-loving p Cork cambium results in the formation becomes impermeable to water accumulation of:	of cork which	These organisms are	fungus like in one phase of oeba like in another phase of (B) slime molds			
37.	 (A) resins (B) suberin (C) lignins (D) tannins The family containing mustard a characters are: (A) Brassicaceae – Tetramerous stamens, bicarpellary gynoecium 	flowers, six	persistent. This residu (A) pericarp (C) chalazosperm	(D) water molds nans of nucellus are also al, persistent nucellus is the (B) perisperm (D) mesosperm			
	fruit (B) Brassicaceae – Pentamerous flestamens, pentacarpellary gynoec	owers, many	(A) Ephedra (C) Pinus	(B) Gnetum (D) Pea			
	type fruit (C) Solanaceae – Pentamerous f stamens, bicarpellary gynoecius	lowers, five	(A) Gracilaris (C) Chondrus	(B) Porphyra(D) Chlorella			
	fruit (D) Poaceae – Trimerous flowers, the monocarpellary gynoecium, cary	nree stamens,	Mesophyll is usually of (A) monocot leaf (C) dorsiventral leaf	(B) isobilateral leaf (D) both (A) and (B)			

fruit

ZOOLOGY

46. The term 'anadromous' in relation to salmons and A competitive inhibitor of an enzyme has which of hilsa pertains to their the following properties? (A) reproduction and excretion (A) It is frequently a feedback inhibitor (B) respiration (B) It becomes covalently attached to an enzyme (C) migration and spawning (C) it interferes with substrate binding to the enzyme (D) maturation of gonads (D) It causes reversible inactivation of the enzyme 47. Match the following host with their parasites 54. Muscle of Boyden and muscle of Oddi are related Taenia A. Pig 2. Liverfluke B. Aedes aegypti (A) pancreas (B) uvula C. Snail (C) stomach (D) intestine 3. Ascaris Functional residual capacity and inspiratory 4. Plague D. Dog capacity when added together give E. None of the five 5. Rabies (A) breathing reserve (B) total lung capacity animals (C) vital capacity 6. Dengue F. Ratflea (D) maximum voluntary ventilation 2 3 4 5 1 6 The correct match is **56.** Е (A) F A В D \mathbf{C} **Blood Groups** Plasma antibodies Е F (B) Α \mathbf{C} D В A. A I. a and b C (C) A В D E F II. b B. B (D) F Е В Α C D C. O III. a 48. In earthworm, the mode of origin of coelom involves D. AB IV. nil. (A) inpushing of ectoderm segmentally, to form C A В D sacs which coalesce to give rise to a continuous (A) I II III IV cavity (B) II Ш I IV IV III II (B) out pocketing of the alimentary canal and the (C) Ι Ш Ι II ΙV pockets coalesce to form a continuous cavity (D) Pick up the correct match 57. (C) separation of mesoderm in the early embryonal A. Oxyntic cells I. Bile period into two layers to create space in between B. Wirsung duct II. HCl C. Brunner's glands III. Pancreatic hormones (D) retention and subsequent enlargement of D. Fat digestion IV. Pancreas embryonal blastocoel to form a space between E. a,b,d cells V. Duodenum mesoderm and endoderm D Α C E 49. Iron free compound in Hb is (A) Ι II III IV V (A) globin (B) haematin (B) II IV V I III (C) bilirubin (D) cyanin (C) V IV III II I 50. The commonest and smallest type of leucocyte in a П I Ш IV (D) healthy adult man is **58.** The total number of lobes and alveoli present in (A) lymphocytes (B) monocytes both the lungs of man is (C) eosinophils (D) basophils (A) 17 and 30 million, respectively (B) 5 and 300 million, respectively 51. The deficiency of the enzyme tyrosinase leads to a (C) 19 and 300 million, respectively serious disorder in the body (D) 18 and 300 lakh, respectively (A) alkaptonuria (B) phenylketonuria **59.** The correct arrangement of leg parts of cockroach (C) tyrosinate (D) albinism One of the following refers to 'feed back (A) coxa, femur, trochanter, tibia and claws mechanism', the effect of (B) coxa, trochanter, femur, tibia, tarsus and claws (A) change in pH and temperature on enzyme (C) coxa, tibia, femur, planulae and claws substrate complex (D) None of the above (B) end product concentration on the rate of 60. The inner layer of blood vessel walls, consisting of enzymatic reactions endothelium, basement membrane, lamina propria

(C) substrate concentration on the rate of

(D) enzymes concentration on the rate of reaction

enzymatic reactions

(B) tunica intima

(D) tunica muscularis

and internal elastic membrane, is the

(A) tunica adventitia

(C) tunica media

ANSWER KEY PHYSICS

1.	(B)	2.	(D)	3.	(D)	4.	(C)	5.	(C)	6.	(D)
7.	(A)	8.	(B)	9.	(D)	10.	(D)	11.	(A)	12.	(B)

13. (B) **14.** (B) **15.** (A)

CHEMISTRY

16.	(A)	17.	(D)	18.	(D)	19.	(C)	20.	(C)	21.	(D)
-----	-----	------------	-----	-----	-----	-----	-----	-----	-----	-----	-----

22. (B) **23.** (B) **24.** (B) **25.** (B) **26.** (A) **27.** (C)

28. (D) **29.** (B) **30.** (B)

BOTANY

31. (B) (A) 34. **36.** (B) 32. 33. (B) (C) **35.** (A) **37.** 40. (A) **38** (C) **39.** (C) (A) 41. (B) **42.** (B)

43. (C) **44.** (D) **45.** (C)

ZOOLOGY

46. (C) **47.** (B) 49. 51. (D) 48. (C) (A) **50.** (A) 52. (B) 53. (C) 54. (A) 55. (B) **56.** (B) 57. (B) **58.** (B) **59.** (B) (B) **60.**