

Postgraduate Programme

Model Question Paper

M.Sc. (Physics)

Time: 2 Hours

Max. Marks : 75

Choose the correct answer and WRITE IN CAPITAL LETTER viz., A, B, C, D or E in the space provided with each question.

SAMPLE QUESTIONS

Section A (25 marks) - 25 questions

1. If the Cartesian coordinates  $(x, y, z)$  of a point are  $(4, 3, 1)$ , its cylindrical polar coordinates  $(\rho, \phi, z)$  will be ( )  
A)  $(5, 36.9^\circ, 1)$   
B)  $(5, 53.1^\circ, 1)$   
C)  $(4, 3, 1)$   
D)  $(\sqrt{7}, 36.9^\circ, 1)$
2.  $V(r, \theta) = \frac{k \cos \theta}{r^2}$ ,  $\nabla V =$  ( )  
A)  $-\frac{k}{r^3}(2 \cos \theta \hat{e}_r + \sin \theta \hat{e}_\theta)$   
B)  $-\frac{k}{r^3}(2 \cos \theta + \sin \theta)$   
C)  $-\frac{k}{r^3} \sqrt{3 \cos^2 \theta + 1}$   
D)  $-\frac{k}{r^3}(2 \cos \theta \hat{e}_r + r \sin \theta \hat{e}_\theta)$
3. The essential singularity of the differential equation:  $x^2(x^2 - 1)y'' + \frac{x}{x+1}y' + \frac{y}{x-1} = 0$  is/are at ( )  
A)  $x = 0$       B)  $x = +1$       C)  $x = -1$       D)  $x = 0, \pm 1$
4. The eigen values of the matrix  $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$  are ( )  
A)  $0, 1$       B)  $+1, -1$       C)  $+i, -i$       D)  $1, i$
5.  $\mathfrak{F}\{f(t - A)\} = e^{i\omega A} F(\omega)$  is called the \_\_\_\_\_ property of the Fourier transformation ( )  
A) Attenuation      B) time Shifting  
C) Convolution      D) Parseval

6. In emitter bias configuration for a transistor,  $V_{BB} = 5V$ ,  $V_{CC} = 15V$ ,  $R_E = 1k\Omega$  and  $R_C = 2k\Omega$ . The collector-emitter voltage  $V_{CE}$  is ( )
- A) 2.1V    B) 4.3V    C) 6.4V    D) 8.6V
7.  $\overline{\overline{A \overline{B \overline{C}}}}$  is equivalent to ( )
- A)  $\overline{A \cdot B \cdot C}$     B)  $\overline{A \cdot \overline{B} \cdot \overline{C}}$     C)  $\overline{A + B + C}$     D)  $A + B + C$
8. Consider the following: Reaction:  $HCl_{(aq)} + NaOH_{(aq)} \rightarrow H_2O_{(l)} + NaCl_{(aq)}$ . The rate of this reaction could be determined by monitoring the change of concentration of: ( )
- A)  $H^+$   
 B)  $Cl^-$   
 C)  $Na^+$   
 D)  $H_2O$
9. Which of the following gives the value stored at the address pointed to by pointer a? ( )
- A) a;    B) val(a);    C) \*a;    D) &a;

## SECTION B – 50 MARKS – (50 QUESTIONS)

### SAMPLE QUESTIONS

1. A particle of mass 100 gm moves on the x-axis under the force field whose potential energy is  $V = \frac{x(x-3)^2}{3}$ . The points of stable equilibrium occur at
- A)  $x = 3$ ,    B)  $x = 1$     C)  $x = 1$  and 3    D) does not exist
2. A particle moves under the influence of the potential  $(x) = A/x^2 - B/x$ . The frequency of small oscillations around the equilibrium point is
- A)  $\sqrt{8mA^3B^4}$     B)  $8mB^4$     C)  $\sqrt{\frac{B^4}{8mA^3}}$     D)  $\sqrt{\frac{8mA^3}{B^4}}$
3. Part of the equation of a plane EM wave travelling in the negative Y direction can be ( )
- A)  $E = A\cos(\omega t - ky)\hat{i}$     B)  $E = A\cos(\omega t - ky)\hat{j}$     C)  $E = A\cos(\omega t + ky)\hat{i}$   
 D)  $E = A\cos(\omega t + ky)\hat{j}$     E)  $E = A\cos(ky - \omega t)\hat{i}$

4. Which of the following equation does not change from one medium to another ( )

- A)  $\vec{\nabla} \cdot \vec{B} = 0$       B)  $\vec{\nabla} \cdot \vec{D} = 0$       C)  $\vec{\nabla}_x \vec{E} = -\partial \vec{B} / \partial t$   
 D)  $\vec{\nabla}_x \vec{H} = \partial \vec{D} / \partial t$       E)  $\vec{\nabla} \cdot \vec{E} = \rho / \epsilon$

5. The ABCD matrix of a thin lens of focal length  $f$  is ( )

- A)  $\begin{pmatrix} 1 & 1/f \\ 0 & 1 \end{pmatrix}$       B)  $\begin{pmatrix} 1 & -1/f \\ 0 & 1 \end{pmatrix}$   
 C)  $\begin{pmatrix} 1 & 0 \\ -1/f & 1 \end{pmatrix}$       D)  $\begin{pmatrix} 1 & 1/f \\ 0 & 1 \end{pmatrix}$       E)  $\begin{pmatrix} -1/f & 1 \\ 0 & 1 \end{pmatrix}$

6. The electric field in a certain region is given by  $\vec{E} = A(yz\hat{i} + xz\hat{k})$ , where  $A = 10 \text{ Nm}^{-2}/\text{C}$  and the potential at the origin of the coordinates is 20 Volts. What will be the potential at a point  $x=2, y=1, z=1$ ? ( all coordinates are in meters) ( )

- A) 10 Volts      B) -30 Volts      C) 30 Volts      D) -10 Volts      E) 0 volts

7. A mathematical approach to the first law of thermo dynamics produced which equation? ( )

- A)  $W + Q = U$       B)  $Q = U + W$       C)  $U = Q - W$   
 D) all the above      E) None of the above


8. A nucleus  $ZXA$  has mass  $M$  kg. If  $M_p$  and  $M_n$  denote the mass (in kg) of proton and neutron respectively, the binding energy in joule is ( )

- A)  $[ZM_p + (A - Z)M_n - M]c^2$       B)  $[ZM_p + ZM_n - M]c^2$   
 C)  $M - ZM_p - (A - Z)M_n$       D)  $[M - ZM_p - (A - Z)M_n]c^2$   
 E)  $A[M_p + M_n]c^2$

9. The eigen values of the operator  $L_z$  are ( )

- A)  $\ell(\ell+1)\hbar^2$       B)  $\ell(\ell+1)\hbar$       C)  $m\hbar$       D)  $m\hbar^2$       E) Zero

10. A mass  $M$  moves with speed  $V$  in the x-direction. It explodes into two pieces that go off at angles  $\theta_1, \theta_2$  as shown in figure. What are the magnitudes of the momenta of the two pieces? ( )



- A)  $p_1 = \frac{P \sin \theta_2}{\sin(\theta_1 + \theta_2)}$ ,  $p_2 = \frac{P \sin \theta_1}{\sin(\theta_1 + \theta_2)}$       B)  $p_1 = \frac{P \sin \theta_2}{\cos(\theta_1 + \theta_2)}$ ,  $p_2 = \frac{P \sin \theta_1}{\cos(\theta_1 + \theta_2)}$   
 C)  $p_1 = \frac{P \sin \theta_2}{\sin(\theta_1 - \theta_2)}$ ,  $p_2 = \frac{P \sin \theta_1}{\sin(\theta_1 - \theta_2)}$       D)  $p_1 = \frac{P \sin \theta_2}{\cos(\theta_1 - \theta_2)}$ ,  $p_2 = \frac{P \sin \theta_1}{\cos(\theta_1 - \theta_2)}$