

Interdisciplinary programme in
INDUSTRIAL ENGINEERING & OPERATIONS RESEARCH
INDIAN INSTITUTE OF TECHNOLOGY BOMBAY

Sample Questions for M.Tech. Admissions Entrance Test
 (some of which appeared in 2006 paper)

Instructions: *No clarifications on the questions should be sought during the examination.
 Calculator not required*

1. The Normal probability distribution is also called _____ (in honor of the person who proposed it as a model for statistical measurement errors)

- (P) Gaussian Distribution (Q) Student's t Distribution
 (R) Bernoulli Distribution (S) Poisson Distribution

2. If the probability of head appearing in a single toss of a coin is p , then the probability that head appears for the first time in the 10th toss is:

- (P) $p(1-p)^9$ (Q) p^{10}
 (R) $p(1-p)$ (S) $(1-p)p^9$

3. A and B working together can finish a job in T days. If A works alone and completes the job, he will take T + 5 days. If B works alone and completes the same job, he will take T + 45 days. What is T?

- (P) 25 (Q) 60 (R) 15 (S) None of these

4. Let random variable X and Y have probability mass as shown in the table on the right side. For example, $P(X=3, Y=0) = 0.2$.

	X			
Y		1	2	3
0		0.1	0.2	0.2
2		0.3	0.2	0

- (i) $P(X = 1) =$ _____
 (ii) $P(X \leq 2 | Y = 0) =$ _____
 (iii) $E[X | Y = 0] =$ _____
 (iv) Let $Z = \min(X, Y)$. Now, $E[Z] =$ _____

5. Pick the random variable with least variance:

- (P) X is Bernoulli with parameter 0.5 (Q) X is uniform over interval [0, 1]
 (R) X is exponential with rate 1 (S) X = 100

6. The maximum value of solution of ordinary differential equation $\frac{d^2 f(x)}{dx^2} = f(x)$


with $f(0) = 1$ is _____.

7. Let X and Y be two random variables with $E[X] = 2$ and $E[Y] = 3$. (For Q 7)
 In the space given on the right side, **plot E[Z]** as a function of α where,

$$Z = \alpha X + \beta Y \text{ for reals } \alpha \text{ and } \beta \text{ such that } \alpha + \beta = 1.$$

8. Let A be a $N \times N$ matrix with each element $\frac{1}{N}$. Now, it is true that:

- (P) Zero is an eigenvalue of A
 (Q) Determinant of A is non-zero
 (R) Determinant of A is zero
 (S) Determinant depends on value of N



9. For any random variable X, it is true that:

- (P) $E[X^2] \geq (E[X])^2$ (Q) $E[X^2] < (E[X])^2$
 (R) $E[X^2] < 0$ and $(E[X])^2 \geq 0$ (S) Such relations depend on the random variable X

10. Find the maxima and/or the minima of the function $f(x) = x^3 + 3x^2 - 24x + 3$.

11. For the following pseudo-code, write the entire output when $n = 10$: (Note: the `write()` function prints the value of its parameters on the screen)

```
x = 0; y = 1;
write(x, y);
while (n != 0)
{
    f = x + y;
    write(f);
    n--;
    x = y;
    y = f;
}
```

12. Consider the Linear Programming model on the right side:

- (i) Identify the solution space using a graph that defines all the feasible solutions of the model.
- (ii) For the given objective, identify the corner point(s) that define the optimum solution.

$$\begin{aligned} \text{Max } z &= 5x + 4y \\ \text{subject to} \\ 6x + 4y &\leq 24 \\ x + 2y &\leq 6 \\ -x + y &\leq 1 \\ y &\leq 2 \\ x, y &\geq 0 \end{aligned}$$

13. For the system of linear equations $Ax = b$ for a square, non-singular matrix A (of dimension n), which of the following are true?

- (P) There is a unique solution of this system for any vector b
- (Q) There is a non zero solution for any non zero vector b
- (R) There is no solution for the case $b = 0$
- (S) There are infinitely many solutions for any vector b

14. The function of two variables $f(x,y) = x^2 - y^2$ over \mathbb{R}^2 has

- (P) A local minimum and a local maximum, but no global minima or maxima
- (Q) No local minimum or local maximum
- (R) No stationary point (where the gradient vector is zero)
- (S) One global minima and one local maxima

15. The transportation problem in linear programming is of the form

$$\begin{aligned} \text{Min } \sum_i \sum_j c_{ij} x_{ij} \\ \text{s.t. } \sum_j x_{ij} &= a_i \text{ for } i \text{ from } 1, \dots, m \\ \sum_i x_{ij} &= b_j \text{ for } j \text{ from } j = 1, \dots, n, \\ \text{all } x_{ij} &\geq 0. \end{aligned}$$

If the transportation problem has an optimal solution, then

- (P) The maximum number of non zero x_{ij} values is m
- (Q) The maximum number of non zero x_{ij} values is n
- (R) The maximum number of non zero x_{ij} values is $m+n$
- (S) The maximum number of non zero x_{ij} values is $m+n-1$

16. With respect to the assignment problem in linear programming, which of the following is true?

- (P) The assignment problem is a special case of the transportation problem
- (Q) The transportation problem is a special case of the assignment problem
- (R) Neither the transportation nor the assignment problems are special cases of the other
- (S) The assignment problem will result in a degenerate solution for the relevant LP

17. The problem: $\text{Max } xyz$ s.t. $x + y + z = 10$, $x, y, z \geq 0$ has

- (P) No feasible solution
- (Q) A unique solution
- (R) Multiple optimal solutions
- (S) Unbounded solution (i.e. no optimal solution)

