

Special Instructions / Useful Data	
\mathbb{R}	Set of all real numbers
\mathbb{R}^n	$\{(x_1, \dots, x_n) : x_i \in \mathbb{R}, i = 1, \dots, n\}$
$P(A)$	Probability of an event A
i.i.d.	Independently and identically distributed
$Bin(n, p)$	Binomial distribution with parameters n and p
$Poisson(\theta)$	Poisson distribution with mean θ
$N(\mu, \sigma^2)$	Normal distribution with mean μ and variance σ^2
$Exp(\lambda)$	The exponential distribution with probability density function $f(x \lambda) = \begin{cases} \lambda e^{-\lambda x}, & x > 0, \\ 0, & \text{otherwise} \end{cases}, \lambda > 0$
t_n	Student's t distribution with n degrees of freedom
χ_n^2	Chi-square distribution with n degrees of freedom
$\chi_{n,\alpha}^2$	A constant such that $P(W > \chi_{n,\alpha}^2) = \alpha$, where W has χ_n^2 distribution
$\Phi(x)$	Cumulative distribution function of $N(0,1)$
$\phi(x)$	Probability density function of $N(0,1)$
A^c	Complement of an event A
$E(X)$	Expectation of a random variable X
$Var(X)$	Variance of a random variable X
$B(m, n)$	$\int_0^1 x^{m-1} (1-x)^{n-1} dx, m > 0, n > 0$
$[x]$	The greatest integer less than or equal to real number x
f'	Derivative of function f
$\Phi(0.25) = 0.5987, \Phi(0.5) = 0.6915, \Phi(0.625) = 0.7341, \Phi(0.71) = 0.7612,$ $\Phi(1) = 0.8413, \Phi(1.125) = 0.8697, \Phi(2) = 0.9772$	

SECTION – A
MULTIPLE CHOICE QUESTIONS (MCQ)

Q. 1 – Q.10 carry one mark each.

Q.1 Let

$$P = \begin{bmatrix} 1 & 2 & 0 & 2 \\ -1 & -2 & 1 & 1 \\ 1 & 2 & -3 & -7 \\ 1 & 2 & -2 & -4 \end{bmatrix}.$$

Then rank of P equals

- (A) 4
- (B) 3
- (C) 2
- (D) 1

Q.2 Let α, β, γ be real numbers such that $\beta \neq 0$ and $\gamma \neq 0$. Suppose

$$P = \begin{bmatrix} \alpha & \beta \\ \gamma & 0 \end{bmatrix},$$

and $P^{-1} = P$. Then

- (A) $\alpha = 0$ and $\beta\gamma = 1$
- (B) $\alpha \neq 0$ and $\beta\gamma = 1$
- (C) $\alpha = 0$ and $\beta\gamma = 2$
- (D) $\alpha = 0$ and $\beta\gamma = -1$

Q.3 Let $m > 1$. The volume of the solid generated by revolving the region between the y -axis and the curve $xy = 4$, $1 \leq y \leq m$, about the y -axis is 15π . The value of m is

- (A) 14
- (B) 15
- (C) 16
- (D) 17

Q.4 Consider the region S enclosed by the surface $z = y^2$ and the planes $z = 1, x = 0, x = 1, y = -1$ and $y = 1$. The volume of S is

- (A) $\frac{1}{3}$
- (B) $\frac{2}{3}$
- (C) 1
- (D) $\frac{4}{3}$

Q.5 Let X be a discrete random variable with the moment generating function

$$M_X(t) = e^{0.5(e^t - 1)}, t \in \mathbb{R}.$$

Then $P(X \leq 1)$ equals

- (A) $e^{-1/2}$ (B) $\frac{3}{2} e^{-1/2}$ (C) $\frac{1}{2} e^{-1/2}$ (D) $e^{-(e-1)/2}$

Q.6 Let E and F be two independent events with

$$P(E|F) + P(F|E) = 1, P(E \cap F) = \frac{2}{9} \text{ and } P(F) < P(E).$$

Then $P(E)$ equals

- (A) $\frac{1}{3}$ (B) $\frac{1}{2}$ (C) $\frac{2}{3}$ (D) $\frac{3}{4}$

Q.7 Let X be a continuous random variable with the probability density function

$$f(x) = \frac{1}{(2+x^2)^{3/2}}, x \in \mathbb{R}.$$

Then $E(X^2)$

- (A) equals 0 (B) equals 1
(C) equals 2 (D) does not exist

Q.8 The probability density function of a random variable X is given by

$$f(x) = \begin{cases} \alpha x^{\alpha-1}, & 0 < x < 1 \\ 0, & \text{otherwise} \end{cases}, \alpha > 0.$$

Then the distribution of the random variable $Y = \log_e X^{-2\alpha}$ is

- (A) χ_2^2 (B) $\frac{1}{2} \chi_2^2$ (C) $2\chi_2^2$ (D) χ_1^2

Q.9 Let X_1, X_2, \dots be a sequence of i.i.d. $N(0,1)$ random variables. Then, as $n \rightarrow \infty$, $\frac{1}{n} \sum_{i=1}^n X_i^2$ converges in probability to

- (A) 0 (B) 0.5 (C) 1 (D) 2

Q.10 Consider the simple linear regression model with n random observations $Y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$, $i = 1, \dots, n$, ($n > 2$). β_0 and β_1 are unknown parameters, x_1, \dots, x_n are observed values of the regressor variable and $\varepsilon_1, \dots, \varepsilon_n$ are error random variables with $E(\varepsilon_i) = 0$, $i = 1, \dots, n$, and for $i, j = 1, \dots, n$, $Cov(\varepsilon_i, \varepsilon_j) = \begin{cases} 0, & \text{if } i \neq j, \\ \sigma^2, & \text{if } i = j. \end{cases}$ For real constants a_1, \dots, a_n , if $\sum_{i=1}^n a_i Y_i$ is an unbiased estimator of β_1 , then

- (A) $\sum_{i=1}^n a_i = 0$ and $\sum_{i=1}^n a_i x_i = 0$ (B) $\sum_{i=1}^n a_i = 0$ and $\sum_{i=1}^n a_i x_i = 1$
 (C) $\sum_{i=1}^n a_i = 1$ and $\sum_{i=1}^n a_i x_i = 0$ (D) $\sum_{i=1}^n a_i = 1$ and $\sum_{i=1}^n a_i x_i = 1$

Q. 11 – Q. 30 carry two marks each.

Q.11 Let (X, Y) have the joint probability density function

$$f(x, y) = \begin{cases} \frac{1}{2} y^2 e^{-x}, & \text{if } 0 < y < x < \infty, \\ 0, & \text{otherwise.} \end{cases}$$

Then $P(Y < 1 | X = 3)$ equals

- (A) $\frac{1}{81}$ (B) $\frac{1}{27}$ (C) $\frac{1}{9}$ (D) $\frac{1}{3}$

Q.12 Let X_1, X_2, \dots be a sequence of i.i.d. random variables having the probability density function

$$f(x) = \begin{cases} \frac{1}{B(6, 4)} x^5 (1-x)^3, & 0 < x < 1, \\ 0, & \text{otherwise.} \end{cases}$$

Let $Y_i = \frac{X_i}{1 - X_i}$ and $U_n = \frac{1}{n} \sum_{i=1}^n Y_i$. If the distribution of $\frac{\sqrt{n}(U_n - 2)}{\alpha}$ converges to $N(0, 1)$ as $n \rightarrow \infty$, then a possible value of α is

- (A) $\sqrt{7}$ (B) $\sqrt{5}$ (C) $\sqrt{3}$ (D) 1

Q.13 Let X_1, \dots, X_n be a random sample from a population with the probability density function

$$f(x|\theta) = \begin{cases} 4e^{-4(x-\theta)}, & x > \theta, \\ 0, & \text{otherwise} \end{cases}, \quad \theta \in \mathbb{R}.$$

If $T_n = \min\{X_1, \dots, X_n\}$, then

- (A) T_n is unbiased and consistent estimator of θ
- (B) T_n is biased and consistent estimator of θ
- (C) T_n is unbiased but NOT consistent estimator of θ
- (D) T_n is NEITHER unbiased NOR consistent estimator of θ

Q.14 Let X_1, \dots, X_n be i.i.d. random variables with the probability density function

$$f(x) = \begin{cases} e^{-x}, & x > 0, \\ 0, & \text{otherwise.} \end{cases}$$

If $X_{(n)} = \max\{X_1, \dots, X_n\}$, then $\lim_{n \rightarrow \infty} P(X_{(n)} - \log_e n \leq 2)$ equals

- (A) $1 - e^{-2}$
- (B) $e^{-e^{-0.5}}$
- (C) $e^{-e^{-2}}$
- (D) e^{-e^2}

Q.15 Let X and Y be two independent $N(0,1)$ random variables. Then $P(0 < X^2 + Y^2 < 4)$ equals

- (A) $1 - e^{-2}$
- (B) $1 - e^{-4}$
- (C) $1 - e^{-1}$
- (D) e^{-2}

Q.16 Let X be a random variable with the cumulative distribution function

$$F(x) = \begin{cases} 0, & x < 0, \\ \frac{x}{8}, & 0 \leq x < 2, \\ \frac{x^2}{16}, & 2 \leq x < 4, \\ 1, & x \geq 4. \end{cases}$$

Then $E(X)$ equals

- (A) $\frac{12}{31}$
- (B) $\frac{13}{12}$
- (C) $\frac{31}{21}$
- (D) $\frac{31}{12}$

Q.17 Let X_1, \dots, X_n be a random sample from a population with the probability density function

$$f(x) = \frac{1}{2\theta} e^{-|x|/\theta}, \quad x \in \mathbb{R}, \theta > 0.$$

For a suitable constant K , the critical region of the most powerful test for testing $H_0 : \theta = 1$ against $H_1 : \theta = 2$ is of the form

- (A) $\sum_{i=1}^n |X_i| > K$ (B) $\sum_{i=1}^n |X_i| < K$
 (C) $\sum_{i=1}^n \frac{1}{|X_i|} < K$ (D) $\sum_{i=1}^n \frac{1}{|X_i|} > K$

Q.18 Let $X_1, \dots, X_n, X_{n+1}, X_{n+2}, \dots, X_{n+m}$ ($n > 4, m > 4$) be a random sample from $N(\mu, \sigma^2)$; $\mu \in \mathbb{R}, \sigma > 0$. If $\bar{X}_1 = \frac{1}{n} \sum_{i=1}^n X_i$ and $\bar{X}_2 = \frac{1}{m-2} \sum_{i=n+1}^{n+m-2} X_i$, then the distribution of the random variable

$$T = \frac{X_{n+m} - X_{n+m-1}}{\sqrt{\sum_{i=1}^n (X_i - \bar{X}_1)^2 + \sum_{i=n+1}^{n+m-2} (X_i - \bar{X}_2)^2}}$$

is

- (A) t_{n+m-2}
 (B) $\sqrt{\frac{2}{n+m-1}} t_{n+m-1}$
 (C) $\sqrt{\frac{2}{n+m-4}} t_{n+m-4}$
 (D) t_{n+m-4}

Q.19 Let X_1, \dots, X_n ($n > 1$) be a random sample from a *Poisson*(θ) population, $\theta > 0$, and

$T = \sum_{i=1}^n X_i$. Then the uniformly minimum variance unbiased estimator of θ^2 is

- (A) $\frac{T(T-1)}{n^2}$ (B) $\frac{T(T-1)}{n(n-1)}$
 (C) $\frac{T(T-1)}{n(n+1)}$ (D) $\frac{T^2}{n^2}$

- Q.20 Let X be a random variable whose probability mass functions $f(x|H_0)$ (under the null hypothesis H_0) and $f(x|H_1)$ (under the alternative hypothesis H_1) are given by

$X = x$	0	1	2	3
$f(x H_0)$	0.4	0.3	0.2	0.1
$f(x H_1)$	0.1	0.2	0.3	0.4

For testing the null hypothesis $H_0: X \sim f(x|H_0)$ against the alternative hypothesis $H_1: X \sim f(x|H_1)$, consider the test given by: Reject H_0 if $X > \frac{3}{2}$.

If $\alpha =$ size of the test and $\beta =$ power of the test, then

- (A) $\alpha = 0.3$ and $\beta = 0.3$
 (B) $\alpha = 0.3$ and $\beta = 0.7$
 (C) $\alpha = 0.7$ and $\beta = 0.3$
 (D) $\alpha = 0.7$ and $\beta = 0.7$
- Q.21 Let X_1, \dots, X_n be a random sample from a $N(2\theta, \theta^2)$ population, $\theta > 0$. A consistent estimator for θ is

- (A) $\frac{1}{n} \sum_{i=1}^n X_i$ (B) $\left(\frac{5}{n} \sum_{i=1}^n X_i^2 \right)^{1/2}$
 (C) $\frac{1}{5n} \sum_{i=1}^n X_i^2$ (D) $\left(\frac{1}{5n} \sum_{i=1}^n X_i^2 \right)^{1/2}$

- Q.22 An institute purchases laptops from either vendor V_1 or vendor V_2 with equal probability. The lifetimes (in years) of laptops from vendor V_1 have a $U(0, 4)$ distribution, and the lifetimes (in years) of laptops from vendor V_2 have an $Exp(1/2)$ distribution. If a randomly selected laptop in the institute has lifetime more than two years, then the probability that it was supplied by vendor V_2 is

- (A) $\frac{2}{2+e}$ (B) $\frac{1}{1+e}$ (C) $\frac{1}{1+e^{-1}}$ (D) $\frac{2}{2+e^{-1}}$

Q.23 Let $y(x)$ be the solution to the differential equation

$$x^4 \frac{dy}{dx} + 4x^3 y + \sin x = 0; \quad y(\pi) = 1, \quad x > 0.$$

Then $y\left(\frac{\pi}{2}\right)$ is

(A) $\frac{10(1+\pi^4)}{\pi^4}$

(B) $\frac{12(1+\pi^4)}{\pi^4}$

(C) $\frac{14(1+\pi^4)}{\pi^4}$

(D) $\frac{16(1+\pi^4)}{\pi^4}$

Q.24 Let $a_n = e^{-2n} \sin n$ and $b_n = e^{-n} n^2 (\sin n)^2$ for $n \geq 1$. Then

(A) $\sum_{n=1}^{\infty} a_n$ converges but $\sum_{n=1}^{\infty} b_n$ does NOT converge

(B) $\sum_{n=1}^{\infty} b_n$ converges but $\sum_{n=1}^{\infty} a_n$ does NOT converge

(C) both $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ converge

(D) NEITHER $\sum_{n=1}^{\infty} a_n$ NOR $\sum_{n=1}^{\infty} b_n$ converges

Q.25 Let

$$f(x) = \begin{cases} x \sin^2(1/x), & x \neq 0, \\ 0, & x = 0, \end{cases} \quad \text{and} \quad g(x) = \begin{cases} x (\sin x) \sin(1/x), & x \neq 0, \\ 0, & x = 0. \end{cases}$$

Then

(A) f is differentiable at 0 but g is NOT differentiable at 0

(B) g is differentiable at 0 but f is NOT differentiable at 0

(C) f and g are both differentiable at 0

(D) NEITHER f NOR g is differentiable at 0

Q.26 Let $f : [0, 4] \rightarrow \mathbb{R}$ be a twice differentiable function. Further, let $f(0) = 1$, $f(2) = 2$ and $f(4) = 3$. Then

- (A) there does NOT exist any $x_1 \in (0, 2)$ such that $f'(x_1) = \frac{1}{2}$
- (B) there exist $x_2 \in (0, 2)$ and $x_3 \in (2, 4)$ such that $f'(x_2) = f'(x_3)$
- (C) $f''(x) > 0$ for all $x \in (0, 4)$
- (D) $f''(x) < 0$ for all $x \in (0, 4)$

Q.27 Let $f(x, y) = x^2 - 400xy^2$ for all $(x, y) \in \mathbb{R}^2$. Then f attains its

- (A) local minimum at $(0, 0)$ but NOT at $(1, 1)$
- (B) local minimum at $(1, 1)$ but NOT at $(0, 0)$
- (C) local minimum both at $(0, 0)$ and $(1, 1)$
- (D) local minimum NEITHER at $(0, 0)$ NOR at $(1, 1)$

Q.28 Let $y(x)$ be the solution to the differential equation

$$4 \frac{d^2 y}{dx^2} + 12 \frac{dy}{dx} + 9y = 0, \quad y(0) = 1, \quad y'(0) = -4.$$

Then $y(1)$ equals

- (A) $-\frac{1}{2} e^{-3/2}$
- (B) $-\frac{3}{2} e^{-3/2}$
- (C) $-\frac{5}{2} e^{-3/2}$
- (D) $-\frac{7}{2} e^{-3/2}$

Q.29 Let $g : [0, 2] \rightarrow \mathbb{R}$ be defined by

$$g(x) = \int_0^x (x-t)e^t dt.$$

The area between the curve $y = g''(x)$ and the x -axis over the interval $[0, 2]$ is

- (A) $e^2 - 1$
- (B) $2(e^2 - 1)$
- (C) $4(e^2 - 1)$
- (D) $8(e^2 - 1)$

Q.30 Let P be a 3×3 singular matrix such that $P\vec{v} = \vec{v}$ for a nonzero vector \vec{v} and

$$P \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = \begin{bmatrix} 2/5 \\ 0 \\ -2/5 \end{bmatrix}.$$

Then

(A) $P^3 = \frac{1}{5}(7P^2 - 2P)$

(B) $P^3 = \frac{1}{4}(7P^2 - 2P)$

(C) $P^3 = \frac{1}{3}(7P^2 - 2P)$

(D) $P^3 = \frac{1}{2}(7P^2 - 2P)$

SECTION - B

MULTIPLE SELECT QUESTIONS (MSQ)

Q. 31 – Q. 40 carry two marks each.

Q.31 For two nonzero real numbers a and b , consider the system of linear equations

$$\begin{bmatrix} a & b \\ b & a \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} b/2 \\ a/2 \end{bmatrix}.$$

Which of the following statements is (are) TRUE?

- (A) If $a = b$, the solutions of the system lie on the line $x + y = 1/2$
- (B) If $a = -b$, the solutions of the system lie on the line $y - x = 1/2$
- (C) If $a \neq \pm b$, the system has no solution
- (D) If $a \neq \pm b$, the system has a unique solution

Q.32 For $n \geq 1$, let

$$a_n = \begin{cases} n 2^{-n}, & \text{if } n \text{ is odd,} \\ -3^{-n}, & \text{if } n \text{ is even.} \end{cases}$$

Which of the following statements is (are) TRUE?

- (A) The sequence $\{a_n\}$ converges
- (B) The sequence $\{|a_n|^{1/n}\}$ converges
- (C) The series $\sum_{n=1}^{\infty} a_n$ converges
- (D) The series $\sum_{n=1}^{\infty} |a_n|$ converges

Q.33 Let $f : (0, \infty) \rightarrow \mathbb{R}$ be defined by

$$f(x) = x \left(e^{1/x^3} - 1 + \frac{1}{x^3} \right).$$

Which of the following statements is (are) TRUE?

- (A) $\lim_{x \rightarrow \infty} f(x)$ exists
- (B) $\lim_{x \rightarrow \infty} x f(x)$ exists
- (C) $\lim_{x \rightarrow \infty} x^2 f(x)$ exists
- (D) There exists $m > 0$ such that $\lim_{x \rightarrow \infty} x^m f(x)$ does NOT exist.

- Q.34 For $x \in \mathbb{R}$, define $f(x) = \cos(\pi x) + [x^2]$ and $g(x) = \sin(\pi x)$. Which of the following statements is (are) TRUE?
- (A) $f(x)$ is continuous at $x = 2$
 (B) $g(x)$ is continuous at $x = 2$
 (C) $f(x) + g(x)$ is continuous at $x = 2$
 (D) $f(x)g(x)$ is continuous at $x = 2$
- Q.35 Let E and F be two events with $0 < P(E) < 1$, $0 < P(F) < 1$ and $P(E|F) > P(E)$. Which of the following statements is (are) TRUE?
- (A) $P(F|E) > P(F)$
 (B) $P(E|F^c) > P(E)$
 (C) $P(F|E^c) < P(F)$
 (D) E and F are independent
- Q.36 Let X_1, \dots, X_n ($n > 1$) be a random sample from a $U(2\theta - 1, 2\theta + 1)$ population, $\theta \in \mathbb{R}$, and $Y_1 = \min\{X_1, \dots, X_n\}$, $Y_n = \max\{X_1, \dots, X_n\}$. Which of the following statistics is (are) maximum likelihood estimator (s) of θ ?
- (A) $\frac{1}{4}(Y_1 + Y_n)$
 (B) $\frac{1}{6}(2Y_1 + Y_n + 1)$
 (C) $\frac{1}{8}(Y_1 + 3Y_n - 2)$
 (D) Every statistic $T(X_1, \dots, X_n)$ satisfying $\frac{(Y_n - 1)}{2} < T(X_1, \dots, X_n) < \frac{(Y_1 + 1)}{2}$
- Q.37 Let X_1, \dots, X_n be a random sample from a $N(0, \sigma^2)$ population, $\sigma > 0$. Which of the following testing problems has (have) the region $\left\{ (x_1, \dots, x_n) \in \mathbb{R}^n : \sum_{i=1}^n x_i^2 \geq \chi_{n, \alpha}^2 \right\}$ as the most powerful critical region of level α ?
- (A) $H_0 : \sigma^2 = 1$ against $H_1 : \sigma^2 = 2$
 (B) $H_0 : \sigma^2 = 1$ against $H_1 : \sigma^2 = 4$
 (C) $H_0 : \sigma^2 = 2$ against $H_1 : \sigma^2 = 1$
 (D) $H_0 : \sigma^2 = 1$ against $H_1 : \sigma^2 = 0.5$

Q.38 Let X_1, \dots, X_n be a random sample from a $N(0, 2\theta^2)$ population, $\theta > 0$. Which of the following statements is (are) TRUE?

- (A) (X_1, \dots, X_n) is sufficient and complete
 (B) (X_1, \dots, X_n) is sufficient but NOT complete
 (C) $\sum_{i=1}^n X_i^2$ is sufficient and complete
 (D) $\frac{1}{2n} \sum_{i=1}^n X_i^2$ is the uniformly minimum variance unbiased estimator for θ^2

Q.39 Let X_1, \dots, X_n be a random sample from a population with the probability density function

$$f(x|\theta) = \begin{cases} \theta e^{-\theta x}, & x > 0, \\ 0, & \text{otherwise} \end{cases}, \theta > 0.$$

Which of the following is (are) $100(1-\alpha)\%$ confidence interval(s) for θ ?

- (A) $\left(\frac{\chi_{2n, 1-\alpha/2}^2}{2 \sum_{i=1}^n X_i}, \frac{\chi_{2n, \alpha/2}^2}{2 \sum_{i=1}^n X_i} \right)$ (B) $\left(0, \frac{\chi_{2n, \alpha}^2}{2 \sum_{i=1}^n X_i} \right)$
 (C) $\left(\frac{\chi_{2n, 1-\alpha/2}^2}{\sum_{i=1}^n X_i}, \frac{\chi_{2n, \alpha/2}^2}{\sum_{i=1}^n X_i} \right)$ (D) $\left(\frac{2 \sum_{i=1}^n X_i}{\chi_{2n, \alpha/2}^2}, \frac{2 \sum_{i=1}^n X_i}{\chi_{2n, 1-\alpha/2}^2} \right)$

Q.40 The cumulative distribution function of a random variable X is given by

$$F(x) = \begin{cases} 0, & x < 2, \\ \frac{1}{10} \left(x^2 - \frac{7}{3} \right), & 2 \leq x < 3, \\ 1, & x \geq 3. \end{cases}$$

Which of the following statements is (are) TRUE?

- (A) $F(x)$ is continuous everywhere
 (B) $F(x)$ increases only by jumps
 (C) $P(X=2) = \frac{1}{6}$
 (D) $P\left(X = \frac{5}{2} \mid 2 \leq X \leq 3\right) = 0$

SECTION – C
NUMERICAL ANSWER TYPE (NAT)

Q. 41 – Q. 50 carry one mark each.

Q.41 Let X_1, \dots, X_{10} be a random sample from a $N(3, 12)$ population. Suppose $Y_1 = \frac{1}{6} \sum_{i=1}^6 X_i$ and $Y_2 = \frac{1}{4} \sum_{i=7}^{10} X_i$. If $\frac{(Y_1 - Y_2)^2}{\alpha}$ has a χ_1^2 distribution, then the value of α is _____

Q.42 Let X be a continuous random variable with the probability density function

$$f(x) = \begin{cases} \frac{2x}{9}, & 0 < x < 3, \\ 0, & \text{otherwise.} \end{cases}$$

Then the upper bound of $P(|X - 2| > 1)$ using Chebyshev's inequality is _____

Q.43 Let X and Y be continuous random variables with the joint probability density function

$$f(x, y) = \begin{cases} e^{-(x+y)}, & -\infty < x, y < 0, \\ 0, & \text{otherwise.} \end{cases}$$

Then $P(X < Y) =$ _____

Q.44 Let X and Y be continuous random variables with the joint probability density function

$$f(x, y) = \frac{1}{2\pi} e^{-(x^2+y^2)/2}, \quad (x, y) \in \mathbb{R}^2.$$

Then $P(X > 0, Y < 0) =$ _____

Q.45 Let Y be a $\text{Bin}\left(72, \frac{1}{3}\right)$ random variable. Using normal approximation to binomial distribution, an approximate value of $P(22 \leq Y \leq 28)$ is _____

Q.46 Let X be a $\text{Bin}(2, p)$ random variable and Y be a $\text{Bin}(4, p)$ random variable, $0 < p < 1$. If

$$P(X \geq 1) = \frac{5}{9}, \text{ then } P(Y \geq 1) = \underline{\hspace{2cm}}$$

Q.47 Consider the linear transformation

$$T(x, y, z) = (2x + y + z, x + z, 3x + 2y + z).$$

The rank of T is $\underline{\hspace{2cm}}$

Q.48 The value of $\lim_{n \rightarrow \infty} n \left[e^{-n} \cos(4n) + \sin\left(\frac{1}{4n}\right) \right]$ is $\underline{\hspace{2cm}}$

Q.49 Let $f: [0, 13] \rightarrow \mathbb{R}$ be defined by $f(x) = x^{13} - e^{-x} + 5x + 6$. The minimum value of the function f on $[0, 13]$ is $\underline{\hspace{2cm}}$

Q.50 Consider a differentiable function f on $[0, 1]$ with the derivative $f'(x) = 2\sqrt{2x}$. The arc length of the curve $y = f(x)$, $0 \leq x \leq 1$, is $\underline{\hspace{2cm}}$

Q. 51 – Q. 60 carry two marks each.

Q.51 Let m be a real number such that $m > 1$. If

$$\int_1^m \int_0^1 \int_{\sqrt{x}}^1 e^{y^3} dy dx dz = e - 1,$$

then $m = \underline{\hspace{2cm}}$

Q.52 Let

$$P = \begin{bmatrix} 1 & -3 & 3 \\ 0 & -5 & 6 \\ 0 & -3 & 4 \end{bmatrix}.$$

The product of the eigen values of P^{-1} is $\underline{\hspace{2cm}}$

Q.53 The value of the real number m in the following equation

$$\int_0^1 \int_x^{\sqrt{2-x^2}} (x^2 + y^2) dy dx = \int_{m\pi}^{\pi/2} \int_0^{\sqrt{2}} r^3 dr d\theta$$

is _____

Q.54 Let $a_1 = 1$ and $a_n = 2 - \frac{1}{n}$ for $n \geq 2$. Then

$$\sum_{n=1}^{\infty} \left(\frac{1}{a_n^2} - \frac{1}{a_{n+1}^2} \right)$$

converges to _____

Q.55 Let X_1, X_2, \dots be a sequence of i.i.d. random variables with the probability density function

$$f(x) = \begin{cases} 4x^2 e^{-2x}, & x > 0, \\ 0, & \text{otherwise} \end{cases}$$

and let $S_n = \sum_{i=1}^n X_i$. Then $\lim_{n \rightarrow \infty} P\left(S_n \leq \frac{3n}{2} + \sqrt{3n}\right)$ is _____

Q.56 Let X and Y be continuous random variables with the joint probability density function

$$f(x, y) = \begin{cases} \frac{c x^2}{y^3}, & 0 < x < 1, y > 1, \\ 0, & \text{otherwise} \end{cases},$$

where c is a suitable constant. Then $E(X) =$ _____

Q.57 Two points are chosen at random on a line segment of length 9 cm. The probability that the distance between these two points is less than 3 cm is _____

Q.58 Let X be a continuous random variable with the probability density function

$$f(x) = \begin{cases} \frac{x+1}{2}, & -1 < x < 1, \\ 0, & \text{otherwise.} \end{cases}$$

Then $P\left(\frac{1}{4} < X^2 < \frac{1}{2}\right) =$ _____

Q.59 If X is a $U(0,1)$ random variable, then $P\left(\min(X, 1-X) \leq \frac{1}{4}\right) =$ _____

Q.60 In a colony all families have at least one child. The probability that a randomly chosen family from this colony has exactly k children is $(0.5)^k$; $k = 1, 2, \dots$. A child is either a male or a female with equal probability. The probability that such a family consists of at least one male child and at least one female child is _____

END OF THE QUESTION PAPER