ACHLA



Test Booklet Code



This Booklet contains 24 pages.

Do not open this Test Booklet until you are asked to do so.

Read carefully the Instructions on the Back Cover of this Test Booklet.

Important Instructions:

- 1. The Answer Sheet is inside this Test Booklet. When you are directed to open the Test Booklet, take out the Answer Sheet and fill in the particulars on **Side-1** and **Side-2** carefully with **blue/black** ball point pen only.
- 2. The test is of **3 hours** duration and this Test Booklet contains **180** questions. Each question carries **4** marks. For each correct response, the candidate will get **4** marks. For each incorrect response, **one mark** will be deducted from the total scores. The maximum marks are 720.
- 3. Use Blue/Black Ball Point Pen only for writing particulars on this page/marking responses.
- 4. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
- 5. On completion of the test, the candidate must hand over the Answer Sheet to the Invigilator before leaving the Room/Hall. The candidates are allowed to take away this Test Booklet with them.
- 6. The CODE for this Booklet is **CC**. Make sure that the CODE printed on **Side-2** of the Answer Sheet is the same as that on this Test Booklet. In case of discrepancy, the candidate should immediately report the matter to the Invigilator for replacement of both the Test Booklet and the Answer Sheet.
- 7. The candidates should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your Roll No. anywhere else except in the specified space in the Test Booklet/Answer Sheet.
- 8. Use of white fluid for correction is **not** permissible on the Answer Sheet.

Name of the Candidate (in Capitals) :	
Roll Number : in figures	
: in words	
Centre of Examination (in Capitals):	
-	Invigilator's Signature :
Facsimile signature stamp of	
Centre Superintendent :	

ACHLA/CC/Page 1 English

1. Niche is

- (1) the functional role played by the organism where it lives
- (2) the range of temperature that the organism needs to live
- (3) the physical space where an organism lives
- (4) all the biological factors in the organism's environment.
- **2.** Which of the following is a secondary pollutant?
 - (1) O_3
 - (2) SO_2
 - (3) CO_2
 - (4) CO
- **3.** In stratosphere, which of the following elements acts as a catalyst in degradation of ozone and release of molecular oxygen?
 - (1) Oxygen
 - (2) Fe
 - (3) Cl
 - (4) Carbon
- 4. World Ozone Day is celebrated on
 - (1) 22nd April
 - (2) 16th September
 - (3) 21st April
 - (4) 5th June
- **5.** What type of ecological pyramid would be obtained with the following data?

Secondary consumer: 120 g

Primary consumer : 60 g

Primary producer : 10 g

- $(1) \quad Upright\ pyramid\ of\ biomass$
- (2) Upright pyramid of numbers
- (3) Pyramid of energy
- (4) Inverted pyramid of biomass
- **6.** Natality refers to
 - (1) Number of individuals entering a habitat
 - (2) Number of individuals leaving the habitat
 - (3) Birth rate
 - (4) Death rate

- 7. Offsets are produced by
 - (1) Parthenogenesis
 - (2) Parthenocarpy
 - (3) Mitotic divisions
 - (4) Meiotic divisions
- **8.** The experimental proof for semiconservative replication of DNA was first shown in a
 - (1) Virus
 - (2) Plant
 - (3) Bacterium
 - (4) Fungus
- **9.** Select the *correct* match :
 - (1) Francois Jacob and Lac operon Jacques Monod
 - (2) Matthew Meselson *Pisum sativum* and F. Stahl
 - (3) Alfred Hershey and TMV Martha Chase
 - $\begin{array}{ccc} \text{(4)} & \text{Alec Jeffreys} & & & Streptococcus \\ & & & pneumoniae \end{array}$
- **10.** Which of the following has proved helpful in preserving pollen as fossils?
 - (1) Sporopollenin
 - (2) Oil content
 - (3) Cellulosic intine
 - (4) Pollenkitt
- **11.** Which of the following pairs is **wrongly** matched?

(1) T.H. Morgan : Linkage

(2) XO type sex : Grasshopper determination

(3) ABO blood grouping : Co-dominance

(4) Starch synthesis in pea : Multiple alleles

- **12.** Which of the following flowers only once in its life-time?
 - (1) Papaya
 - (2) Mango
 - (3) Jackfruit
 - (4) Bamboo species
- **13.** Select the *correct* statement :
 - (1) Transduction was discovered by S. Altman.
 - (2) Spliceosomes take part in translation.
 - (3) Punnett square was developed by a British scientist.
 - (4) Franklin Stahl coined the term "linkage".

	(3) NADH		(4)	i	iv	iii	ii	
	(3) NADH		(4)	i	iv	iii	ii	
	(2) NADPH		(3)	iii	ii	i	iv	
	• 0							
	(1) Oxygen		(2)		iv	iii	i	
	(1) Oxygen		(2)	ii	iv	iii	i	
			(2)	ii	iv	iii	i	
					iv	iii	i	
	reaction of photosynthesis?		(1)		iv	i	ii	
			(1)	iii	iv	i	ii	
			(1)	iii	iv	i	ii	
			(1)	iii	iv	i	ii	
			(1)	iii	iv	i	ii	
			(1)	iii	iv	i	ii	
18.	Which of the following is not a product of light reaction of photosynthesis?		(1)	iii	iv	i	ii	
			(1)	iii	iv	i	ii	
			(1)	iii	iv	i	ii	
			(1)	iii	iv	i	ii	
			(1)	iii	iv	i	ii	
			(1)	iii	iv	i	ii	
			(1)	iii	iv	i	ii	
	reaction of photosynthesis?		(1)	iii	iv	i	ii	
	reaction of photosynthesis?		(1)		1V	1	11	
			(2)	ii	iv	iii	i	
	(1) Oxygen		(2)	ii	iv	iii	i	
	(1) Oxygen		(2)	11	1V	111	1	
	• 0							
	(2) NADPH		(3)	iii	ii	i	iv	
	(3) NADH		(4)	i	iv	iii	ii	
			(4)	1	1V	111	11	
			\ - /	-				
	(4) ATP			_		_		
10		23.	Whi	ch on	e is <i>wr</i>	nglv	matched?	
19.	Which of the following is true for nucleolus?	zs.						
10.	_		(1)	Unio	cellular	orgar	nism – <i>Chlorella</i>	;
	(1) It is a site for active ribosomal RNA							
			(2)	Gen	ıma cup	os	- Marchan	tia
	synthesis.				_			
	(2) It takes part in spindle formation		(3)	Bifla	agellate	zoosp	ores - Brown al	gae
	(2) It takes part in spindle formation.				_	_		_
			(4)	Unit	lagella	te gan	netes – <i>Polysiphe</i>	onia
	(2) It is a mambrana based attended	1			-	_		
	(3) It is a membrane-bound structure.						11	
	(4) Larger nucleoli are present in dividing cells.	24.	Afte	r kar	yogamy	follo	wed by meiosis, spo	res are
20.	(4) Larger nucleoli are present in dividing cells.	24.					· -	res are
20.	(4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote?	24.		duced	exogen	ously	· -	res are
20.	(4) Larger nucleoli are present in dividing cells.	24.	proc	duced	exogen	ously	· -	ores are
20.	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria 	24.	prod (1)	duced Saco	exogen	ously	· -	ores are
20.	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria (2) Nostoc 	24.	proc	duced	exogen	ously	· -	ores are
20.	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria 	24.	prod (1) (2)	duced Saco Agai	exogen charom; ricus	ously	· -	ores are
20.	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria (2) Nostoc (3) Mycobacterium 	24.	prod (1) (2) (3)	duced Saco Agai Alte	exogen charom ricus rnaria	ously yces	· -	ores are
20.	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria (2) Nostoc 	24.	prod (1) (2)	duced Saco Agai Alte	exogen charom; ricus	ously yces	· -	ores are
	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria (2) Nostoc (3) Mycobacterium (4) Saccharomyces 	24.	prod (1) (2) (3)	duced Saco Agai Alte	exogen charom ricus rnaria	ously yces	· -	ores are
20. 21.	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria (2) Nostoc (3) Mycobacterium (4) Saccharomyces The two functional groups characteristic of 		prod (1) (2) (3) (4)	Sace Agai Alte Neu	exogen charomy ricus rnaria rospora	ously yces	in	ores are
	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria (2) Nostoc (3) Mycobacterium (4) Saccharomyces 	24. 25.	prod (1) (2) (3) (4)	Sace Agai Alte Neu	exogen charomy ricus rnaria rospora	ously yces	· -	ores are
	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria (2) Nostoc (3) Mycobacterium (4) Saccharomyces The two functional groups characteristic of sugars are 		prod (1) (2) (3) (4) Win	Sace Agai Alte Neu	exogen charomy ricus rnaria rospora ollen gr	ously yces	in	ores are
	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria (2) Nostoc (3) Mycobacterium (4) Saccharomyces The two functional groups characteristic of sugars are (1) carbonyl and hydroxyl 		prod (1) (2) (3) (4) Win (1)	Sacci Agai Alter Neu ged p	exogen charomy ricus rnaria rospora ollen gr	ously yces	in	ores are
	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria (2) Nostoc (3) Mycobacterium (4) Saccharomyces The two functional groups characteristic of sugars are 		prod (1) (2) (3) (4) Win	Sace Agai Alte Neu	exogen charomy ricus rnaria rospora ollen gr	ously yces	in	ores are
	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria (2) Nostoc (3) Mycobacterium (4) Saccharomyces The two functional groups characteristic of sugars are (1) carbonyl and hydroxyl (2) carbonyl and phosphate 		proc (1) (2) (3) (4) Win (1) (2)	Agar Alter Neu ged po Pinu Man	exogen charomy ricus rnaria rospora ollen gr us	ously yces	in	ores are
	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria (2) Nostoc (3) Mycobacterium (4) Saccharomyces The two functional groups characteristic of sugars are (1) carbonyl and hydroxyl 		proc (1) (2) (3) (4) Win (1) (2) (3)	Agar Agar Alter Neu ged po Pini Man	exogen charomy ricus rnaria rospora ollen gr us us	ously yces	in	ores are
	 (4) Larger nucleoli are present in dividing cells. Which among the following is <i>not</i> a prokaryote? (1) Oscillatoria (2) Nostoc (3) Mycobacterium (4) Saccharomyces The two functional groups characteristic of sugars are (1) carbonyl and hydroxyl (2) carbonyl and phosphate 		proc (1) (2) (3) (4) Win (1) (2)	Agar Agar Alter Neu ged po Pini Man	exogen charomy ricus rnaria rospora ollen gr us	ously yces	in	ores are

- **26.** Which of the following is commonly used as a vector for introducing a DNA fragment in human lymphocytes?
 - (1) pBR 322
 - (2) λ phage
 - (3) Ti plasmid
 - (4) Retrovirus
- **27.** A 'new' variety of rice was patented by a foreign company, though such varieties have been present in India for a long time. This is related to
 - (1) Basmati
 - (2) Lerma Rojo
 - (3) Sharbati Sonora
 - (4) Co-667
- 28. Use of bioresources by multinational companies and organisations without authorisation from the concerned country and its people is called
 - (1) Bioexploitation
 - (2) Biodegradation
 - (3) Biopiracy
 - (4) Bio-infringement
- **29.** Select the *correct* match :
 - (1) G. Mendel Transformation
 - (2) T.H. Morgan Transduction
 - (3) $F_2 \times \text{Recessive parent}$ Dihybrid cross
 - (4) Ribozyme Nucleic acid
- **30.** The correct order of steps in Polymerase Chain Reaction (PCR) is
 - (1) Denaturation, Annealing, Extension
 - (2) Denaturation, Extension, Annealing
 - (3) Annealing, Extension, Denaturation
 - (4) Extension, Denaturation, Annealing
- **31.** In India, the organisation responsible for assessing the safety of introducing genetically modified organisms for public use is
 - (1) Genetic Engineering Appraisal Committee (GEAC)
 - (2) Research Committee on Genetic Manipulation (RCGM)
 - (3) Council for Scientific and Industrial Research (CSIR)
 - (4) Indian Council of Medical Research (ICMR)

- **32.** What is the role of NAD⁺ in cellular respiration?
 - (1) It is the final electron acceptor for anaerobic respiration.
 - (2) It is a nucleotide source for ATP synthesis.
 - (3) It functions as an electron carrier.
 - (4) It functions as an enzyme.
- **33.** Which one of the following plants shows a very close relationship with a species of moth, where none of the two can complete its life cycle without the other?
 - (1) Viola
 - (2) Banana
 - (3) Yucca
 - (4) Hydrilla
- **34.** Pollen grains can be stored for several years in liquid nitrogen having a temperature of
 - $(1) 160^{\circ}C$
 - $(2) 196^{\circ}C$
 - $(3) 80^{\circ}C$
 - $(4) 120^{\circ}C$
- **35.** In which of the following forms is iron absorbed by plants?
 - (1) Both ferric and ferrous
 - (2) Free element
 - (3) Ferrous
 - (4) Ferric
- **36.** Double fertilization is
 - (1) Syngamy and triple fusion
 - (2) Fusion of two male gametes with one egg
 - (3) Fusion of one male gamete with two polar nuclei
 - (4) Fusion of two male gametes of a pollen tube with two different eggs
- **37.** Oxygen is *not* produced during photosynthesis by
 - (1) Chara
 - (2) Cycas
 - (3) Nostoc
 - (4) Green sulphur bacteria
- **38.** Which of the following elements is responsible for maintaining turgor in cells?
 - (1) Calcium
 - (2) Potassium
 - (3) Sodium
 - (4) Magnesium

- **39.** Pneumatophores occur in
 - (1) Submerged hydrophytes
 - (2) Carnivorous plants
 - (3) Free-floating hydrophytes
 - (4) Halophytes

40. Select the **wrong** statement :

- (1) Mitochondria are the powerhouse of the cell in all kingdoms except Monera.
- (2) Pseudopodia are locomotory and feeding structures in Sporozoans.
- (3) Mushrooms belong to Basidiomycetes.
- (4) Cell wall is present in members of Fungi and Plantae.
- **41.** Secondary xylem and phloem in dicot stem are produced by
 - (1) Axillary meristems
 - (2) Phellogen
 - (3) Vascular cambium
 - (4) Apical meristems
- **42.** Sweet potato is a modified
 - (1) Rhizome
 - (2) Tap root
 - (3) Adventitious root
 - (4) Stem
- **43.** Which of the following statements is *correct*?
 - (1) Stems are usually unbranched in both *Cycas* and *Cedrus*.
 - (2) Horsetails are gymnosperms.
 - (3) Selaginella is heterosporous, while Salvinia | 50. is homosporous.
 - (4) Ovules are not enclosed by ovary wall in gymnosperms.
- 44. Casparian strips occur in
 - (1) Endodermis
 - (2) Cortex
 - (3) Pericycle
 - (4) Epidermis
- **45.** Plants having little or no secondary growth are
 - (1) Cycads
 - (2) Conifers
 - (3) Deciduous angiosperms
 - (4) Grasses

- 46. Nissl bodies are mainly composed of
 - (1) Free ribosomes and RER
 - (2) Nucleic acids and SER
 - (3) DNA and RNA
 - (4) Proteins and lipids
- **47.** Which of these statements is *incorrect*?
 - (1) Oxidative phosphorylation takes place in outer mitochondrial membrane.
 - (2) Glycolysis operates as long as it is supplied with NAD that can pick up hydrogen atoms.
 - (3) Glycolysis occurs in cytosol.
 - (4) Enzymes of TCA cycle are present in mitochondrial matrix.
- 48. Many ribosomes may associate with a single mRNA to form multiple copies of a polypeptide simultaneously. Such strings of ribosomes are termed as
 - (1) Nucleosome
 - (2) Plastidome
 - (3) Polyhedral bodies
 - (4) Polysome
- **49.** Which of the following terms describe human dentition?
 - (1) Pleurodont, Diphyodont, Heterodont
 - (2) Pleurodont, Monophyodont, Homodont
 - (3) Thecodont, Diphyodont, Heterodont
 - (4) Thecodont, Diphyodont, Homodont
- **50.** Which of the following events does **not** occur in rough endoplasmic reticulum?
 - (1) Phospholipid synthesis
 - (2) Cleavage of signal peptide
 - (3) Protein glycosylation
 - (4) Protein folding
- **51.** Select the *incorrect* match :
 - (1) Polytene Oocytes of amphibians chromosomes
 - (2) Submetacentric L-shaped chromosomes chromosomes
 - (3) Allosomes Sex chromosomes
 - (4) Lampbrush Diplotene bivalents

- **52.** Which of the following is an amino acid derived hormone?
 - (1) Estriol
 - (2) Estradiol
 - (3) Ecdysone
 - (4) Epinephrine
- **53.** Which of the following structures or regions is *incorrectly* paired with its function?

(1) Corpus callosum : band of fibers

connecting left and right cerebral hemispheres.

(2) Hypothalamus : production of

releasing hormones and regulation of temperature,

hunger and thirst.

(3) Limbic system : consists of fibre

tracts that interconnect different regions of

brain; controls movement.

(4) Medulla oblongata: controls respiration

and cardiovascular

reflexes.

- **54.** Which of the following hormones can play a significant role in osteoporosis?
 - (1) Parathyroid hormone and Prolactin
 - (2) Estrogen and Parathyroid hormone
 - (3) Progesterone and Aldosterone
 - (4) Aldosterone and Prolactin
- **55.** The transparent lens in the human eye is held in its place by
 - (1) smooth muscles attached to the ciliary body
 - (2) smooth muscles attached to the iris
 - (3) ligaments attached to the iris
 - (4) ligaments attached to the ciliary body

- **56.** In a growing population of a country,
 - (1) pre-reproductive individuals are less than the reproductive individuals.
 - (2) reproductive and pre-reproductive individuals are equal in number.
 - (3) reproductive individuals are less than the post-reproductive individuals.
 - (4) pre-reproductive individuals are more than the reproductive individuals.
- **57.** Match the items given in Column I with those in Column II and select the *correct* option given below:

 $Column\ I$

Column II

- a. Eutrophication i. UV-B radiation
- b. Sanitary landfill ii. Deforestation
- c. Snow blindness
- iii. Nutrient

enrichment

d. Jhum cultivation iv. Waste disposal

	a	b	\mathbf{c}	d
(1)	i	ii	iv	iii
(2)	iii	iv	i	ii
(3)	i	iii	iv	ii

i

58. Which part of poppy plant is used to obtain the drug "Smack"?

iii

iv

(1) Leaves

ii

(2) Roots

(4)

- (3) Latex
- (4) Flowers
- **59.** Which one of the following population interactions is widely used in medical science for the production of antibiotics?
 - (1) Amensalism
 - (2) Parasitism
 - (3) Mutualism
 - (4) Commensalism
- **60.** All of the following are included in 'Ex-situ conservation' *except*
 - (1) Seed banks
 - (2) Botanical gardens
 - (3) Sacred groves
 - (4) Wildlife safari parks

61.	· · · · · · · · · · · · · · · · · · ·					65.	AGGTATCGCAT is a sequence from the coding strand of a gene. What will be the corresponding					
	help in erythropoiesis?						sequence of the transcribed mRNA?					
	(1)		etal cell	S			(1)	UCC	CAUAG	CGUA		
	(2)	Gob	let cells				(2)	ACC	UAUG	CGAU		
	(3)	Mucous cells Chief cells					(3)	UGO	GTUTC	GCAT		
	(4)						(4)	AGC	HUAUC	GCAU		
62.	2. Match the items given in Column I with those in Column II and select the <i>correct</i> option given below:				66.	X chromosomes. This chromosome of inherited by						
		$Column\ I$			$Column\ II$		(1) Both sons and daughters(2) Only grandchildren					5
	a.	Fibr	inogen	i.	Osmotic balance		(3)	-	sons	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	. -	
	b.	Glok	oulin	ii.	Blood clotting		(4)	-	daugh	ters		
	c.	Albı	ımin	iii.	Defence mechanism	67.	Match the items given in Column I with those in					umn I with those in
		_	L.				Column II and select the <i>correct</i> option given					
	(1)	a ii	b iii	c i			belo		_			
	(2)		iii	ii					mn I			Column II
	(3)	i i	ii	iii			a.	Prol	iferative	e Phase	e i.	Breakdown of endometrial
	(4)	iii	ii	i								lining
	(1)						b.	Secr	etory P	hase	ii.	Follicular Phase
63.		Calcium is important in skeletal muscle					c.	Men	struatio	n	iii.	Luteal Phase
	cont		n becau					a	b	c		
	(1)	_			ion of bonds between		(1)	iii	i	ii		
		the myosin cross bridges and the actin filament.					(2)	ii	iii	i		
	(2)			e mvosir	n head from the actin		(3)	i	iii	ii		
	(-)	filament.		1 11000 110111 0110 000111		(4)	iii	ii	i			
	(3)	activates the myosin ATPase by binding to it.			ATPase by binding to	68.		ording lution		go de	Vries	, the mechanism of
	(4)	bind	ls to tro	ponin to	remove the masking of		(1)	Min	or muta	tions		
		activ	ve sites o	on actin f	or myosin.		(2)	Phe	notypic	variatio	ons	
64.	Whi	ch o	f the	following	is an occupational		(3)		ation			
			ry disord	_			(4)	Mul	tiple ste	p muta	tions	i
	(1)	Emp	ohysema	l		69.				ıg are p	art o	f an operon <i>except</i>
	(2) Botulism					(1)	a pr	omoter				

(3)

(4)

Silicosis

Anthracis

(2)

(3)

(4)

an enhancer

an operator

structural genes

- 70. Which of the following options correctly represents the lung conditions in asthma and emphysema, respectively?
 - (1) Decreased respiratory surface; Inflammation of bronchioles
 - (2) Increased respiratory surface; Inflammation of bronchioles
 - (3) Increased number of bronchioles; Increased respiratory surface
 - (4) Inflammation of bronchioles; Decreased respiratory surface
- **71.** Match the items given in Column I with those in Column II and select the *correct* option given below:

$Column\ I$

Column II

- a. Tricuspid valve i
- . Between left atrium and left ventricle
- b. Bicuspid valve
- ii. Between right ventricle and pulmonary artery
- c. Semilunar valve iii. Between right atrium and right ventricle
- $\begin{array}{cccc} \mathbf{a} & \mathbf{b} & \mathbf{c} \\ \text{(1)} & \text{ii} & \text{i} & \text{iii} \end{array}$
- $(2) \quad i \qquad \quad ii \qquad \quad iii$
- (3) i iii ii
- (4) iii i ii
- **72.** Match the items given in Column I with those in Column II and select the *correct* option given below:

Column I

Column II


- a. Tidal volume
- i. 2500 3000 mL
- b. Inspiratory Reserve volume
- ii. 1100 1200 mL
- c. Expiratory Reserve
- iii. 500 550 mL
- d. Residual volume
- iv. 1000 1100 mL
- a b c d
- (1) iv iii ii i
- (2) i iv ii iii
- (3) iii i iv ii
- (4) iii ii iv


- **73.** Hormones secreted by the placenta to maintain pregnancy are
 - (1) hCG, progestogens, estrogens, glucocorticoids
 - (2) hCG, hPL, progestogens, estrogens
 - (3) hCG, hPL, estrogens, relaxin, oxytocin
 - (4) hCG, hPL, progestogens, prolactin
- **74.** The contraceptive 'SAHELI'
 - (1) is a post-coital contraceptive.
 - (2) is an IUD.
 - (3) increases the concentration of estrogen and prevents ovulation in females.
 - (4) blocks estrogen receptors in the uterus, preventing eggs from getting implanted.
- **75.** The difference between spermiogenesis and spermiation is
 - (1) In spermiogenesis spermatozoa are formed, while in spermiation spermatozoa are released from sertoli cells into the cavity of seminiferous tubules.
 - (2) In spermiogenesis spermatozoa from sertoli cells are released into the cavity of seminiferous tubules, while in spermiation spermatozoa are formed.
 - (3) In spermiogenesis spermatozoa are formed, while in spermiation spermatids are formed.
 - (4) In spermiogenesis spermatids are formed, while in spermiation spermatozoa are formed.
- **76.** The amnion of mammalian embryo is derived from
 - (1) ectoderm and endoderm
 - (2) mesoderm and trophoblast
 - $(3) \quad endoderm \ and \ mesoderm$
 - (4) ectoderm and mesoderm


77.		ch of the following animals does <i>not</i> undergo amorphosis?	83.	Match the items given in Column I with those in Column II and select the <i>correct</i> option given							
	(1)	Starfish		belo							
	(2)	Moth			Colu	mn~I		Col	lumn II		
	(3)	Tunicate									
	(4)	Earthworm		a.	Glyc	osuria	i.		amulation of uric in joints		
78.	Whi hom	ch one of these animals is not a leotherm?		b.	Gout ii.			s of crystallised s within the kidney			
	(1)	Psittacula		c.	Rena	al calculi	iii.	Infla	ammation in		
	(2)	Camelus						glomeruli			
	(3)	Chelone		d.	Glomerular nephritis		iv.	Presence of glucose in urine			
	(4)	Macropus			_						
79.	Whi	ch of the following features is used to identify		(4)	a	b	c	d 			
		ale cockroach from a female cockroach?		(1)	iv	i	ii	ii	i		
	(1)	Presence of anal cerci		(2)	ii	iii	i	iv	V		
	(2)	Forewings with darker tegmina		(3)	i	ii	iii	iv	V		
	(3)	Presence of caudal styles		(4)	iii	ii	iv	i			
	(4)	Presence of a boat shaped sternum on the 9 th abdominal segment	84.	Match the items given in Column II and select th					n Column I with those in he <i>correct</i> option given		
80.	Which of the following organisms are known as chief producers in the oceans?			below:							
	(1)	Euglenoids				mn I			Column~II		
	(2)	Cyanobacteria			(Function)				(Part of Excretory		
	(3)	Diatoms							System)		
	(4)	Dinoflagellates		a.	Ultra	afiltratio	n	i.	Henle's loop		
81.	Cilia	ates differ from all other protozoans in		b.	Conc of ur	entration	n	ii.	Ureter		
	(1)	having two types of nuclei		c.	Tran	sport of		iii.	Urinary bladder		
	(2)	using pseudopodia for capturing prey			urine			·			
	(3)	having a contractile vacuole for removing excess water		d.				Malpighian corpuscle			
	(4)	using flagella for locomotion						v.	Proximal		
82.	Iden	tify the vertebrate group of animals						٧.	convoluted tubule		
02.	chai	characterized by crop and gizzard in its digestive system.			a	b	c	d	1		
	(1)	Osteichthyes		(1)	v	iv	i	ii	i		
	(2)	Aves		(2)	v	iv	i	ii	Ĺ		
	(3)	Reptilia		(3)	iv	i	ii	ii	i		
	(4)	Amphibia		(4)	iv	v	ii	ii			
				-							


- **85.** Among the following sets of examples for divergent evolution, select the *incorrect* option :
 - (1) Eye of octopus, bat and man
 - (2) Brain of bat, man and cheetah
 - (3) Heart of bat, man and cheetah
 - (4) Forelimbs of man, bat and cheetah
- **86.** Conversion of milk to curd improves it nutritional value by increasing the amount of
 - (1) Vitamin E
 - (2) Vitamin B_{12}
 - (3) Vitamin A
 - (4) Vitamin D
- **87.** Which of the following characteristics represent 'Inheritance of blood groups' in humans?
 - a. Dominance
 - b. Co-dominance
 - c. Multiple allele
 - d. Incomplete dominance
 - e. Polygenic inheritance
 - (1) a, c and e
 - (2) b, d and e
 - (3) a, b and c
 - (4) b, c and e
- **88.** Which of the following is *not* an autoimmune disease?
 - (1) Vitiligo
 - (2) Alzheimer's disease
 - (3) Rheumatoid arthritis
 - (4) Psoriasis
- **89.** The similarity of bone structure in the forelimbs of many vertebrates is an example of
 - (1) Adaptive radiation
 - (2) Convergent evolution
 - (3) Analogy
 - (4) Homology
- **90.** In which disease does mosquito transmitted pathogen cause chronic inflammation of lymphatic vessels?
 - (1) Amoebiasis
 - (2) Ringworm disease
 - (3) Ascariasis
 - (4) Elephantiasis

- 91. A carbon resistor of (47 ± 4.7) k Ω is to be marked with rings of different colours for its identification. The colour code sequence will be
 - (1) Green Orange Violet Gold
 - (2) Yellow Green Violet Gold
 - $(3) \quad Yellow-\ Violet-Orange-Silver$
 - (4) Violet Yellow Orange Silver
- **92.** A set of 'n' equal resistors, of value 'R' each, are connected in series to a battery of emf 'E' and internal resistance 'R'. The current drawn is I. Now, the 'n' resistors are connected in parallel to the same battery. Then the current drawn from battery becomes 10 I. The value of 'n' is
 - (1) 9
 - (2) 20
 - (3) 11
 - (4) 10
- **93.** A battery consists of a variable number 'n' of identical cells (having internal resistance 'r' each) which are connected in series. The terminals of the battery are short-circuited and the current I is measured. Which of the graphs shows the correct relationship between I and n?










- The power radiated by a black body is P and it 94. radiates maximum energy at wavelength, λ_0 . If the temperature of the black body is now changed so that it radiates maximum energy at wavelength $\frac{3}{4}\lambda_0$, the power radiated by it becomes nP. The value of n is
 - **(1)** 256
 - 256 (2)81
 - (3)
 - (4)
- 95. have the same volume. The first wire has cross-sectional area A and the second wire has cross-sectional area 3A. If the length of the first wire is increased by Δl on applying a force F, how much force is needed to stretch the second wire by the same amount?
 - F (1)
 - (2)4 F
 - (3)6 F
 - (4)9 F
- 96. A sample of 0.1 g of water at 100°C and normal pressure $(1.013 \times 10^5 \text{ Nm}^{-2})$ requires 54 cal of heat energy to convert to steam at 100°C. If the volume of the steam produced is 167·1 cc, the change in internal energy of the sample, is
 - (1) 84.5 J
 - (2)42.2 J
 - 208·7 J (3)
 - (4)104·3 J
- 97. A small sphere of radius 'r' falls from rest in a viscous liquid. As a result, heat is produced due to viscous force. The rate of production of heat when the sphere attains its terminal velocity, is proportional to
 - (1) r^4
 - (2)
 - (3)
 - (4)

- The moment of the force, $\overrightarrow{F} = 4\hat{i} + 5\hat{j} 6\hat{k}$ at (2, 0, -3), about the point (2, -2, -2), is given by
 - $(1) -7\hat{i} -4\hat{j} -8\hat{k}$
 - (2) $-7\hat{i} 8\hat{j} 4\hat{k}$
 - $(3) \quad -4\stackrel{\land}{i} \stackrel{\land}{j} 8\stackrel{\land}{k}$
 - $(4) 8\hat{i} 4\hat{i} 7\hat{k}$
- 99. A student measured the diameter of a small steel ball using a screw gauge of least count 0.001 cm. The main scale reading is 5 mm and zero of circular scale division coincides with 25 divisions above the reference level. If screw gauge has a zero error of -0.004 cm, the correct diameter of the ball is
 - **(1)** 0.529 cm
 - (2)0.053 cm
 - 0.525 cm (3)
 - (4) 0.521 cm
- Two wires are made of the same material and | 100. A toy car with charge q moves on a frictionless horizontal plane surface under the influence of a uniform electric field \vec{E} . Due to the force $q\vec{E}$, its velocity increases from 0 to 6 m/s in one second duration. At that instant the direction of the field is reversed. The car continues to move for two more seconds under the influence of this field. The average velocity and the average speed of the toy car between 0 to 3 seconds are respectively
 - 1.5 m/s, 3 m/s(1)
 - (2)1 m/s, 3.5 m/s
 - (3)1 m/s, 3 m/s
 - 2 m/s, 4 m/s(4)
 - **101.** A block of mass m is placed on a smooth inclined wedge ABC of inclination θ as shown in the figure. The wedge is given an acceleration 'a' towards the right. The relation between a and θ for the block to remain stationary on the wedge is




- **(1)** $a = g \tan \theta$
- (2) $a = g \cos \theta$
- $a = \frac{g}{\sin \theta}$
- (4)

- 102. An em wave is propagating in a medium with a velocity $\vec{V} = V\, \hat{i}$. The instantaneous oscillating electric field of this em wave is along +y axis. Then the direction of oscillating magnetic field of the em wave will be along
 - (1) x direction
 - (2) v direction
 - (3) + z direction
 - (4) z direction
- 103. The refractive index of the material of a prism is $\sqrt{2}$ and the angle of the prism is 30°. One of the two refracting surfaces of the prism is made a mirror inwards, by silver coating. A beam of monochromatic light entering the prism from the other face will retrace its path (after reflection from the silvered surface) if its angle of incidence on the prism is
 - (1) zero
 - (2) 30°
 - (3) 45°
 - (4) 60°
- **104.** The magnetic potential energy stored in a certain inductor is 25 mJ, when the current in the inductor is 60 mA. This inductor is of inductance
 - (1) 13.89 H
 - (2) 1.389 H
 - (3) 138·88 H
 - $(4) \quad 0.138 \text{ H}$
- **105.** An object is placed at a distance of 40 cm from a concave mirror of focal length 15 cm. If the object is displaced through a distance of 20 cm towards the mirror, the displacement of the image will be
 - (1) 36 cm towards the mirror
 - (2) 30 cm towards the mirror
 - (3) 36 cm away from the mirror
 - (4) 30 cm away from the mirror

- **106.** The ratio of kinetic energy to the total energy of an electron in a Bohr orbit of the hydrogen atom, is
 - (1) 1:-2
 - (2) 2:-1
 - (3) 1:-1
 - (4) 1:1
- 107. An electron of mass m with an initial velocity $\vec{V} = \overset{\wedge}{V_0} \hat{i} \ (V_0 > 0) \quad \text{enters} \quad \text{an electric field}$ $\vec{E} = \overset{\wedge}{E_0} \hat{i} \ (E_0 = \text{constant} > 0) \text{ at } t = 0. \text{ If } \lambda_0 \text{ is}$ its de-Broglie wavelength initially, then its de-Broglie wavelength at time t is
 - (1) λ_0
 - (2) λ_0 t
 - $(3) \quad \lambda_0 \left(1 + \frac{eE_0}{mV_0} t \right)$
 - $(4) \qquad \frac{\lambda_0}{\left(1+\frac{eE_0}{mV_0}t\right)}$
- 108. When the light of frequency $2\nu_0$ (where ν_0 is threshold frequency), is incident on a metal plate, the maximum velocity of electrons emitted is v_1 . When the frequency of the incident radiation is increased to $5\nu_0$, the maximum velocity of electrons emitted from the same plate is v_2 . The ratio of v_1 to v_2 is
 - (1) 2:1
 - (2) 4:1
 - (3) 1:4
 - (4) 1:2
- 109. For a radioactive material, half-life is 10 minutes. If initially there are 600 number of nuclei, the time taken (in minutes) for the disintegration of 450 nuclei is
 - (1) 15
 - (2) 30
 - (3) 10
 - (4) 20

110. The volume (V) of a monatomic gas varies with its temperature (T), as shown in the graph. The ratio of work done by the gas, to the heat absorbed by it, when it undergoes a change from state A to state B, is



- $(1) \quad \frac{2}{7}$
- (2) $\frac{1}{3}$
- (3) $\frac{2}{3}$
- $(4) \frac{2}{5}$
- 111. The fundamental frequency in an open organ pipe is equal to the third harmonic of a closed organ pipe. If the length of the closed organ pipe is 20 cm, the length of the open organ pipe is
 - (1) 16 cm
 - (2) 12·5 cm
 - (3) 8 cm
 - (4) 13.2 cm
- 112. The efficiency of an ideal heat engine working between the freezing point and boiling point of water, is
 - (1) 12.5%
 - $(2) \quad 6.25\%$
 - (3) 20%
 - (4) 26.8%
- **113.** At what temperature will the rms speed of oxygen molecules become just sufficient for escaping from the Earth's atmosphere?


 (Given:

Mass of oxygen molecule (m) = 2.76×10^{-26} kg Boltzmann's constant $k_B = 1.38 \times 10^{-23}$ J K⁻¹)

- (1) $1.254 \times 10^4 \text{ K}$
- (2) $5.016 \times 10^4 \text{ K}$
- (3) $8.360 \times 10^4 \text{ K}$
- (4) $2.508 \times 10^4 \text{ K}$

- 114. Unpolarised light is incident from air on a plane surface of a material of refractive index ' μ '. At a particular angle of incidence 'i', it is found that the reflected and refracted rays are perpendicular to each other. Which of the following options is correct for this situation ?
 - $(1) \quad i = tan^{-1} \left(\frac{1}{\mu}\right)$
 - $(2) \quad i = \sin^{-1} \left(\frac{1}{\mu}\right)$
 - (3) Reflected light is polarised with its electric vector perpendicular to the plane of incidence
 - (4) Reflected light is polarised with its electric vector parallel to the plane of incidence
- 115. In Young's double slit experiment the separation d between the slits is 2 mm, the wavelength λ of the light used is 5896 Å and distance D between the screen and slits is 100 cm. It is found that the angular width of the fringes is 0·20°. To increase the fringe angular width to 0·21° (with same λ and D) the separation between the slits needs to be changed to
 - (1) 1.7 mm
 - (2) 2.1 mm
 - (3) 1.9 mm
 - (4) 1.8 mm
- **116.** An astronomical refracting telescope will have large angular magnification and high angular resolution, when it has an objective lens of
 - (1) small focal length and small diameter
 - (2) large focal length and large diameter
 - (3) large focal length and small diameter
 - (4) small focal length and large diameter


117. In the circuit shown in the figure, the input voltage V_i is 20 V, V_{BE} = 0 and V_{CE} = 0. The values of I_B , I_C and β are given by



- (1) $I_B = 40 \mu A$, $I_C = 5 mA$, $\beta = 125$
- (2) $I_B = 20 \mu A$, $I_C = 5 mA$, $\beta = 250$
- (3) $I_B = 25 \mu A$, $I_C = 5 mA$, $\beta = 200$
- $(4) \hspace{0.5cm} I_B = 40 \hspace{0.1cm} \mu A, \hspace{0.1cm} I_C = 10 \hspace{0.1cm} mA, \hspace{0.1cm} \beta = 250 \hspace{0.1cm}$

118. In a p-n junction diode, change in temperature due to heating

- (1) affects the overall V I characteristics of p-n junction
- (2) does not affect resistance of p-n junction
- (3) affects only forward resistance
- (4) affects only reverse resistance
- 119. In the combination of the following gates the output Y can be written in terms of inputs A and B as




- (1) $\overline{A+B}$
- (2) $\overline{A \cdot B} + A \cdot B$
- (3) $A \cdot \overline{B} + \overline{A} \cdot B$
- (4) $\overline{A \cdot B}$

- 20. A metallic rod of mass per unit length 0.5 kg m⁻¹ is lying horizontally on a smooth inclined plane which makes an angle of 30° with the horizontal. The rod is not allowed to slide down by flowing a current through it when a magnetic field of induction 0.25 T is acting on it in the vertical direction. The current flowing in the rod to keep it stationary is
 - (1) 11·32 A
 - (2) 14.76 A
 - $(3) \quad 5.98 \text{ A}$
 - (4) 7.14 A
- 121. An inductor 20 mH, a capacitor 100 μF and a resistor 50 Ω are connected in series across a source of emf, V = 10 sin 314 t. The power loss in the circuit is
 - (1) 1·13 W
 - (2) 2·74 W
 - (3) 0·43 W
 - (4) 0.79 W
- 122. A thin diamagnetic rod is placed vertically between the poles of an electromagnet. When the current in the electromagnet is switched on, then the diamagnetic rod is pushed up, out of the horizontal magnetic field. Hence the rod gains gravitational potential energy. The work required to do this comes from
 - (1) the induced electric field due to the changing magnetic field
 - (2) the lattice structure of the material of the rod
 - (3) the magnetic field
 - (4) the current source
- **123.** Current sensitivity of a moving coil galvanometer is 5 div/mA and its voltage sensitivity (angular deflection per unit voltage applied) is 20 div/V. The resistance of the galvanometer is
 - (1) 500Ω
 - (2) 250Ω
 - (3) 25 Ω
 - (4) 40Ω


- 124. A tuning fork is used to produce resonance in a 128. The kinetic energies of a planet in an elliptical glass tube. The length of the air column in this tube can be adjusted by a variable piston. At room temperature of 27°C two successive resonances are produced at 20 cm and 73 cm of column length. If the frequency of the tuning fork is 320 Hz, the velocity of sound in air at 27°C is
 - 300 m/s **(1)**
 - (2)350 m/s
 - (3)339 m/s
 - (4)330 m/s
- **125.** The electrostatic force between the metal plates of an isolated parallel plate capacitor C having a charge Q and area A, is
 - inversely proportional to the distance between the plates.
 - (2)proportional to the square root of the distance between the plates.
 - (3)linearly proportional to the distance between the plates.
 - (4)independent of the distance between the plates.
- **126.** A pendulum is hung from the roof of a sufficiently high building and is moving freely to and fro like a simple harmonic oscillator. The acceleration of the bob of the pendulum is 20 m/s² at a distance of 5 m from the mean position. The time period of oscillation is
 - (1) $1 \mathrm{s}$
 - (2) $2 \mathrm{s}$
 - (3) πs
 - (4) $2\pi s$
- 127. An electron falls from rest through a vertical distance h in a uniform and vertically upward directed electric field E. The direction of electric field is now reversed, keeping its magnitude the same. A proton is allowed to fall from rest in it through the same vertical distance h. The time of fall of the electron, in comparison to the time of fall of the proton is
 - (1) egual
 - (2)10 times greater
 - (3)5 times greater
 - (4)smaller

orbit about the Sun, at positions A, B and C are K_A , K_B and K_C , respectively. AC is the major axis and SB is perpendicular to AC at the position of the Sun S as shown in the figure. Then



- $K_{\rm R} > K_{\rm A} > K_{\rm C}$ (1)
- (2) $K_R < K_\Delta < K_C$
- (3) $K_{\Delta} > K_{B} > K_{C}$
- $K_A < K_B < K_C$ (4)
- 129. A solid sphere is in rolling motion. In rolling motion a body possesses translational kinetic energy (K_t) as well as rotational kinetic energy (K_r) simultaneously. The ratio $K_t : (K_t + K_r)$ for the sphere is
 - (1) 2:5
 - (2)10:7
 - (3)5:7
 - (4) 7:10
- 130. If the mass of the Sun were ten times smaller and the universal gravitational constant were ten times larger in magnitude, which of the following is *not* correct?
 - 'g' on the Earth will not change. **(1)**
 - (2)Time period of a simple pendulum on the Earth would decrease.
 - (3)Walking on the ground would become more difficult.
 - (4)Raindrops will fall faster.
- **131.** A solid sphere is rotating freely about its symmetry axis in free space. The radius of the sphere is increased keeping its mass same. Which of the following physical quantities would remain constant for the sphere?
 - (1) Angular momentum
 - (2)Rotational kinetic energy
 - (3)Moment of inertia
 - Angular velocity (4)

132. A body initially at rest and sliding along a 136. Iron carbonyl, Fe(CO)₅ is frictionless track from a height h (as shown in the figure) just completes a vertical circle of diameter AB = D. The height h is equal to



- (1)
- (2)
- (3)D
- (4)
- 133. Three objects, A: (a solid sphere), B: (a thin circular disk) and C: (a circular ring), each have the same mass M and radius R. They all spin with the same angular speed ω about their own symmetry axes. The amounts of work (W) required to bring them to rest, would satisfy the relation
 - $(1) \quad W_{A} > W_{C} > W_{R}$
 - $(2) W_{B} > W_{A} > W_{C}$
 - (3) $W_A > W_B > W_C$
 - $(4) W_C > W_B > W_A$
- 134. Which one of the following statements is incorrect?
 - Coefficient **(1)** ofsliding friction has dimensions of length.
 - (2)Frictional force opposes the relative motion.
 - (3)Limiting value of static friction is directly proportional to normal reaction.
 - (4) Rolling friction is smaller than sliding friction.
- 135. A moving block having mass m, collides with another stationary block having mass 4m. The lighter block comes to rest after collision. When the initial velocity of the lighter block is v, then the value of coefficient of restitution (e) will be
 - (1) 0.4
 - (2)0.8
 - (3)0.25
 - (4) 0.5


- - (1) dinuclear
 - (2)trinuclear
 - (3)mononuclear
 - tetranuclear (4)
- **137.** Match the metal ions given in Column I with the spin magnetic moments of the ions given in Column II and assign the *correct* code:

	_		
	$Column\ I$		$Column \ II$
a.	Co^{3+}	i.	$\sqrt{8}$ B.M.
b.	Cr^{3+}	ii.	$\sqrt{35}$ B.M.
c.	Fe^{3+}	iii.	$\sqrt{3}$ B.M.
d.	Ni^{2+}	iv.	$\sqrt{24}$ B.M.
		v.	$\sqrt{15}$ B.M.

	a	b	\mathbf{c}	d
(1)	iii	v	i	ii
(2)	iv	i	ii	iii
(3)	i	ii	iii	iv
(4)		**	::	:

- 138. Which one of the following ions exhibits d-d transition and paramagnetism as well?
 - MnO_4^{2-} (1)
 - MnO_4 (2)
 - $\operatorname{Cr}_2\operatorname{O}_7^{2-}$ (3)
 - $\operatorname{CrO}_4^{2-}$
- 139. The geometry and magnetic behaviour of the complex [Ni(CO)₄] are
 - tetrahedral geometry and paramagnetic (1)
 - (2)square planar geometry and paramagnetic
 - (3)tetrahedral geometry and diamagnetic
 - (4) square planar geometry and diamagnetic
- **140.** The type of isomerism shown by the complex $[CoCl_2(en)_2]$ is
 - **(1)** Linkage isomerism
 - (2)Ionization isomerism
 - (3)Coordination isomerism
 - (4)Geometrical isomerism

141. Identify the major products P, Q and R in the 143. For the redox reaction following sequence of reactions:



P Q R

$$(1) \begin{picture}(1){c} CH(CH_3)_2\\ , \begin{picture}(2){c} CH_3 - CO - CH_3\\ \end{picture}$$

$$(2) \quad \bigcirc \overset{\mathrm{CH}(\mathrm{CH}_3)_2}{,} \quad \bigcirc \overset{\mathrm{OH}}{\longrightarrow} , \quad \mathrm{CH_3CH}(\mathrm{OH})\mathrm{CH_3}$$

(4)
$$CH_2CH_2CH_3$$
 CHO , $CH_3CH_2 - OH$

- 142. Which of the following compounds can form a zwitterion?
 - (1) Glycine
 - Benzoic acid (2)
 - (3)Acetanilide
 - Aniline (4)

$$\operatorname{MnO}_4^- + \operatorname{C_2O_4^{2-}} + \operatorname{H}^+ {\longrightarrow} \operatorname{Mn}^{2+} + \operatorname{CO_2} + \operatorname{H_2O}$$

the correct coefficients of the reactants for the balanced equation are


	MnO_4^-	$C_2^{O_4^{2}}$	$\mathrm{H}^{ au}$
(1)	5	16	2
(2)	2	16	5
(3)	2	5	16


- (4)2
- **144.** The correction factor 'a' to the ideal gas equation corresponds to
 - forces of **(1)** attraction between molecules
 - (2)electric field present between the molecules
 - (3)volume of the gas molecules
 - (4)density of the gas molecules
- **145.** Which one of the following conditions will favour maximum formation of the product in the reaction,


$$A_{2}(g) + B_{2}(g) \rightleftharpoons X_{2}(g) \quad \Delta_{r}H = -X kJ$$
?


- (1) High temperature and low pressure
- (2)High temperature and high pressure
- (3)Low temperature and low pressure
- (4)Low temperature and high pressure
- **146.** The bond dissociation energies of X_2 , Y_2 and XYare in the ratio of 1:0.5:1. ΔH for the formation of XY is -200 kJ mol⁻¹. The bond dissociation energy of X2 will be
 - 400 kJ mol^{-1} (1)
 - 800 kJ mol^{-1} (2)
 - 100 kJ mol^{-1} (3)
 - 200 kJ mol^{-1} (4)
- **147.** When initial concentration of the reactant is doubled, the half-life period of a zero order reaction
 - (1) remains unchanged
 - (2)is tripled
 - (3)is doubled
 - (4)is halved

- **148.** Which of the following molecules represents the order of hybridisation sp², sp², sp, sp from left to right atoms?
 - (1) $CH_3 CH = CH CH_3$
 - (2) $CH_2 = CH CH = CH_2$
 - (3) $CH_2 = CH C \equiv CH$
 - (4) $HC \equiv C C \equiv CH$
- **149.** Which of the following carbocations is expected to be most stable?









- 150. Which of the following is correct with respect toI effect of the substituents ? (R = alkyl)
 - (1) $-NR_2 > -OR > -F$
 - $(2) \quad -\mathrm{NH}_2>-\mathrm{OR}>-\mathrm{F}$
 - $(3) \quad -NR_2 < -OR < -F$
 - $(4) NH_2 < -OR < -F$

- **151.** The correct difference between first- and second-order reactions is that
 - (1) the rate of a first-order reaction does depend on reactant concentrations; the rate of a second-order reaction does not depend on reactant concentrations
 - (2) a first-order reaction can be catalyzed; a second-order reaction cannot be catalyzed
 - (3) the half-life of a first-order reaction does not depend on [A]₀; the half-life of a second-order reaction does depend on [A]₀
 - (4) the rate of a first-order reaction does not depend on reactant concentrations; the rate of a second-order reaction does depend on reactant concentrations
- **152.** Among CaH₂, BeH₂, BaH₂, the order of ionic character is
 - (1) BaH₂ < BeH₂ < CaH₂
 - $(2) \quad \text{BeH}_2 < \text{BaH}_2 < \text{CaH}_2$
 - (3) CaH₂ < BeH₂ < BaH₂
 - $(4) \quad \text{BeH}_2 < \text{CaH}_2 < \text{BaH}_2$
- **153.** Consider the change in oxidation state of Bromine corresponding to different emf values as shown in the diagram below:

$$BrO_4^- \xrightarrow{1.82 \text{ V}} BrO_3^- \xrightarrow{1.5 \text{ V}} HBrO$$

$$Br^- \xleftarrow{1.0652 \text{ V}} Br_2 \xleftarrow{1.595 \text{ V}}$$

Then the species undergoing disproportionation is

- (1) HBrO
- (2) Br_2
- (3) BrO_4^-
- (4) BrO $_3^-$
- **154.** In which case is the number of molecules of water maximum?
 - (1) 10^{-3} mol of water
 - (2) 0.00224 L of water vapours at 1 atm and 273 K
 - (3) 0.18 g of water
 - (4) 18 mL of water

- 155. The compound A on treatment with Na gives B, and with PCl₅ gives C. B and C react together to give diethyl ether. A, B and C are in the order
 - (1) C₂H₅OH, C₂H₅ONa, C₂H₅Cl
 - (2)C₂H₅Cl, C₂H₆, C₂H₅OH
 - C₂H₅OH, C₂H₅Cl, C₂H₅ONa (3)
 - C₂H₅OH, C₂H₆, C₂H₅Cl (4)
- **156.** Hydrocarbon (A) reacts with bromine substitution to form an alkyl bromide which by Wurtz reaction is converted to hydrocarbon containing less than four carbon atoms. (A) is
 - (1) CH_{4}
 - (2) $CH_3 - CH_3$
 - $CH_2 = CH_2$ (3)
 - $CH \equiv CH$ (4)
- **157.** The compound C_7H_8 undergoes the following reactions:

$$C_7H_8 \xrightarrow{3 Cl_2/\Delta} A \xrightarrow{Br_2/Fe} B \xrightarrow{Zn/HCl} C$$

The product 'C' is

- (1) *p*-bromotoluene
- (2)3-bromo-2,4,6-trichlorotoluene
- o-bromotoluene (3)
- *m*-bromotoluene (4)
- 158. Which oxide of nitrogen is **not** a common | 163. Regarding cross-linked or network polymers, pollutant introduced into the atmosphere both due to natural and human activity?
 - (1) NO
 - (2) N_2O
 - (3) NO_2
 - **(4)** N_2O_5

- **159.** A mixture of 2·3 g formic acid and 4·5 g oxalic acid is treated with conc. H₂SO₄. The evolved gaseous mixture is passed through KOH pellets. Weight (in g) of the remaining product at STP will be
 - **(1)** 4.4
 - (2)2.8
 - (3)3.0
 - (4) 1.4
- **160.** The difference between amylose and amylopectin
 - Amylose is made up of glucose **(1)** galactose
 - (2)Amylopectin have $1 \rightarrow 4$ α -linkage and $1 \rightarrow 6 \beta$ -linkage
 - (3) $1 \rightarrow 4$ Amvlose have α-linkage and $1 \rightarrow 6 \beta$ -linkage
 - Amylopectin have $1 \rightarrow 4$ α -linkage and $1 \rightarrow 6 \alpha$ -linkage
- 161. Which of the following oxides is most acidic in nature?
 - (1) CaO
 - (2)BaO
 - (3)BeO
 - (4) MgO
- 162. Nitration of aniline in strong acidic medium also gives m-nitroaniline because
 - In acidic (strong) medium aniline is present as anilinium ion.
 - (2)In absence of substituents nitro group always goes to m-position.
 - (3)electrophilic substitution reactions amino group is meta directive.
 - (4) In spite of substituents nitro group always goes to only m-position.
- which of the following statements is *incorrect*?
 - They contain strong covalent bonds in their polymer chains.
 - (2)Examples are bakelite and melamine.
 - (3)They are formed from bi- and tri-functional monomers.
 - (4) They contain covalent bonds between various linear polymer chains.

- 164. Following solutions were prepared by mixing different volumes of NaOH and HCl of different concentrations:
 - a. $60 \text{ mL } \frac{\text{M}}{10} \text{ HCl} + 40 \text{ mL } \frac{\text{M}}{10} \text{ NaOH}$
 - b. 55 mL $\frac{M}{10}$ HCl + 45 mL $\frac{M}{10}$ NaOH
 - c. 75 mL $\frac{M}{5}$ HCl + 25 mL $\frac{M}{5}$ NaOH
 - d. 100 mL $\frac{M}{10}$ HCl + 100 mL $\frac{M}{10}$ NaOH

pH of which one of them will be equal to 1?

- (1) c
- (2) d
- (3) a
- (4) b
- **165.** On which of the following properties does the coagulating power of an ion depend?
 - (1) The sign of charge on the ion alone
 - (2) Both magnitude and sign of the charge on the ion
 - (3) Size of the ion alone
 - (4) The magnitude of the charge on the ion alone
- **166.** The solubility of $BaSO_4$ in water is $2\cdot42\times10^{-3}~gL^{-1}$ at 298 K. The value of its solubility product (K_{sp}) will be

(Given molar mass of $BaSO_4 = 233 \text{ g mol}^{-1}$)

- (1) $1.08 \times 10^{-8} \text{ mol}^2 \text{ L}^{-2}$
- (2) $1.08 \times 10^{-14} \text{ mol}^2 \text{ L}^{-2}$
- (3) $1.08 \times 10^{-12} \text{ mol}^2 \text{ L}^{-2}$
- (4) $1.08 \times 10^{-10} \text{ mol}^2 \text{ L}^{-2}$
- **167.** Given van der Waals constant for NH_3 , H_2 , O_2 and CO_2 are respectively 4·17, 0·244, 1·36 and 3·59, which one of the following gases is most easily liquefied?
 - (1) CO₂
 - (2) O_2
 - (3) H_2
 - (4) NH₃

- 168. Magnesium reacts with an element (X) to form an ionic compound. If the ground state electronic configuration of (X) is 1s² 2s² 2p³, the simplest formula for this compound is
 - $(1) \quad Mg_3X_2$
 - (2) Mg_2X
 - (3) MgX₂
 - $(4) Mg_2X_3$
- **169.** Iron exhibits bcc structure at room temperature. Above 900°C, it transforms to fcc structure. The ratio of density of iron at room temperature to that at 900°C (assuming molar mass and atomic radii of iron remains constant with temperature) is
 - $(1) \quad \frac{1}{2}$
 - $(2) \qquad \frac{3\sqrt{3}}{4\sqrt{2}}$
 - $(3) \quad \frac{4\sqrt{3}}{3\sqrt{2}}$
 - $(4) \qquad \frac{\sqrt{3}}{\sqrt{2}}$
- is 170. Which one is a **wrong** statement?
 - (1) The value of m for d_{z^2} is zero.
 - (2) The electronic configuration of N atom is

$$\begin{array}{ccc}
1s^2 & 2s^2 & 2p_x^1 & 2p_y^1 & 2p_z^1 \\
\uparrow \downarrow & \uparrow \downarrow & \uparrow \downarrow
\end{array}$$

- (3) An orbital is designated by three quantum numbers while an electron in an atom is designated by four quantum numbers.
- (4) Total orbital angular momentum of electron in 's' orbital is equal to zero.
- **171.** Consider the following species:

CN+, CN-, NO and CN

Which one of these will have the highest bond order?

- (1) CN
- (2) CN^+
- (3) CN^-
- (4) NO

- 172. Which of the following statements is **not** true for 178. In the reaction halogens?
 - **(1)** Chlorine has the highest electron-gain enthalpy.
 - (2)All but fluorine show positive oxidation states.
 - (3)All are oxidizing agents.
 - All form monobasic oxyacids. (4)
- 173. Which one of the following elements is unable to form MF_6^{3-} ion?
 - (1) In
 - (2)В
 - (3)Al
 - (4) Ga
- 174. In the structure of ClF₃, the number of lone pairs of electrons on central atom 'Cl' is
 - (1) three
 - (2)four
 - (3)two
 - (4) one
- 175. Considering Ellingham diagram, which of the following metals can be used to reduce alumina?
 - (1) Cu
 - (2)Mg
 - (3)Zn
 - (4)Fe
- 176. The correct order of atomic radii in group 13 elements is
 - B < Ga < Al < In < Tl(1)
 - B < Ga < Al < Tl < In(2)
 - (3)B < Al < Ga < In < Tl
 - B < Al < In < Ga < Tl(4)
- 177. The correct order of N-compounds in its decreasing order of oxidation states is
 - NH₄Cl, N₂, NO, HNO₃
 - HNO₃, NH₄Cl, NO, N₂ (2)
 - HNO₃, NO, NH₄Cl, N₂
 - HNO₃, NO, N₂, NH₄Cl

the electrophile involved is

- (1) dichlorocarbene (:CCl₂)
- dichloromethyl anion (CHCl₂) (2)
- (3)formyl cation (CHO)
- (4) dichloromethyl cation (CHCl₂)
- 179. Carboxylic acids have higher boiling points than aldehydes, ketones and even alcohols comparable molecular mass. It is due to their
 - formation of intermolecular H-bonding
 - more extensive association of carboxylic (2)acid via van der Waals force of attraction
 - (3) formation of carboxylate ion
 - formation of intramolecular H-bonding (4)
- 180. Compound A, C₈H₁₀O, is found to react with NaOI (produced by reacting Y with NaOH) and yields a yellow precipitate with characteristic smell.

A and Y are respectively

(2)
$$\sim$$
 CH – CH $_3$ and I $_2$ OH

(3)
$$\langle \text{CH}_2 - \text{CH}_2 - \text{OH} \text{ and } \text{I}_2 \rangle$$

(4)
$$H_3C \longrightarrow CH_2 - OH \text{ and } I_2$$

SPACE FOR ROUGH WORK

ACHLA/CC/Page 22 English

SPACE FOR ROUGH WORK

ACHLA/CC/Page 23 English

Read carefully the following instructions:

- 1. Each candidate must show on demand his/her Admit Card to the Invigilator.
- 2. No candidate, without special permission of the Superintendent or Invigilator, would leave his/her seat.
- 3. The candidates should not leave the Examination Hall without handing over their Answer Sheet to the Invigilator on duty and sign the Attendance Sheet twice. Cases where a candidate has not signed the Attendance Sheet second time will be deemed not to have handed over the Answer Sheet and dealt with as an unfair means case.
- 4. Use of Electronic/Manual Calculator is prohibited.
- 5. The candidates are governed by all Rules and Regulations of the examination with regard to their conduct in the Examination Hall. All cases of unfair means will be dealt with as per Rules and Regulations of this examination.
- 6. No part of the Test Booklet and Answer Sheet shall be detached under any circumstances.
- 7. The candidates will write the Correct Test Booklet Code as given in the Test Booklet/Answer Sheet in the Attendance Sheet.

ACHLA/CC/Page 24 English